



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>



QB 28 219

YC 15051

LIBRARY
UNIVERSITY OF CALIFORNIA

ST. LOUIS EXHIBIT

NO. 76

LIBRARY
OF THE
UNIVERSITY OF CALIFORNIA.
GIFT OF
Arthur S. Gable
Class 984m
E11

MINERAL TABLES

FOR THE DETERMINATION OF MINERALS BY
THEIR PHYSICAL PROPERTIES.


BY

ARTHUR S. EAKLE, P.H.D.,

Assistant Professor of Mineralogy, University of California,

FIRST EDITION.

FIRST THOUSAND.



NEW YORK:

JOHN WILEY & SONS.

LONDON: CHAPMAN & HALL, LIMITED.

1904.

Copyright, 1904,
BY
ARTHUR S. EAKLE.

ROBERT DRUMMOND, PRINTER, NEW YORK.

INTRODUCTION.

THE natural method of identifying minerals is by using those characters which are prominent or which can be determined in the field or laboratory in the simplest manner. Practice in the determination of minerals by their physical properties tends to develop the habit of close and careful observation, and at the same time enables the student to acquire more knowledge of minerals in a given time than could be obtained by any other method. Experience has demonstrated that work in blowpipe analysis is less apt to become merely mechanical if it has been preceded by such practice.

The tables include the common minerals and a few others of local prominence, which are generally considered as rare in occurrence. The minerals are arranged primarily according to streak and color, as seen in the Analytical Key, and under each color the arrangement is according to hardness. The tables differ from those of Weisbach chiefly in disregarding luster as an important division and in maintaining the same system of arrangement throughout. Various works on mineralogy, especially Dana's System of Mineralogy, have been consulted in the preparation of the tables. For valuable suggestions and criticisms the author is especially indebted to Professor Charles Palache of Harvard, who used the manuscript copies of the tables in the Summer School of the University of California.

BERKELEY, August 25, 1903.

iii

MINERAL TABLES.

PHYSICAL PROPERTIES.



THE only apparatus needed for the tables is a pocket-knife, a horse-shoe magnet, a pocket-lens, a piece of unglazed porcelain or streak-plate, and a scale of hardness consisting of nine minerals. This scale can be procured of any mineral dealer or can be made up from specimens in an ordinary collection of minerals.

Color.—The color of a mineral is perhaps the most important property used in its identification, yet it is the most difficult to accurately describe. A mineral may have but one characteristic color, or it may occur in various colors and shades; consequently a mineral may be repeated several times in the tables. The color must always be judged by the fresh surface or fracture, and it should be homogeneous throughout the mineral. Vitreous minerals may often be discolored or stained by impurities, when they manifestly belong to the colorless or white class.

Luster.—The luster of a mineral refers to the kind of reflected surface. The common kinds are: *metallic*; *submetallic*=imperfect metallic; *vitreous*=glassy; *adamantine*=brilliant oily luster; *resinous*; *greasy*; *pearly*; *silky*; *waxy*.

Streak.—The streak of a mineral is the color of its fine powder. It is best obtained by rubbing the specimen upon the streak-plate until a definite color is produced.

Hardness.—By hardness is meant the resistance that a mineral offers to abrasion or scratching. The relative hardness of a mineral is usually determined by scratching it successively by minerals or substances of known hardness, two minerals of the same hardness mutually scratching each other. The scale of hardness in common use, called the Moh's scale, in ascending degree of hardness, is as follows: 1, Talc; 2, Gypsum; 3, Calcite; 4, Fluorite; 5, Apatite; 6, Feldspar; 7, Quartz; 8, Topaz; 9, Corundum; 10, Diamond.

The thumb-nail will scratch minerals up to 2½, and the ordinary knife-blade up to 5½ in the scale; with a little practice the relative hardness of a mineral under 6 can be approximately determined with a knife-blade. Above 6 the scale of hardness is necessary. The relative hardness of many of the metallic or submetallic minerals can often be judged by the ease or difficulty in obtaining a streak on the streak-plate. Fine fibrous and fine granular minerals usually appear to be much softer than the individual fiber or grain would be, if it were coarser. Also the surface of some minerals is often much softer than the fresh interior, owing to alteration.

Specific Gravity.—The gravity of a mineral is its weight compared with the weight of an equal volume of water. It is determined by first weighing the mineral in air and then weighing it suspended in water.

If w = weight in air, and w' = weight in water, then $G = \frac{w}{w - w'}$. The gravity of minerals can be determined with a chemical balance or

with the convenient Jolly spring-balance. Whether the mineral is light, medium, heavy, or very heavy can often be judged simply by hefting it.

Crystallization.—A few mineral substances on assuming a solid condition do not crystallize and are said to be amorphous, but most mineral substances when solidifying have the property of crystallizing into certain definite crystal forms, and any such crystalline substance must belong to one of six different crystal systems. These systems are: 1, *Isometric*; 2, *Tetragonal*; 3, *Hexagonal*; 4, *Orthorhombic*; 5, *Monoclinic*; 6, *Triclinic*.

Assuming that each system has axes which are intersected by the crystal planes, then:

1. *Isometric* has three equal axes all at right angles. The common forms in this system are: *Cube*, having six square faces; *Octahedron*, having eight equilateral triangular faces; *Rhombic dodecahedron*, having twelve rhombic faces; *Icosatetrahedron* or *Trapezohedron*, having twenty-four trapezohedral faces; *Tetrahedron*, having four equilateral triangular faces; *Pentagonal dodecahedron* or *Pyritohedron*, having twelve pentagonal faces.
2. *Tetragonal* has two horizontal axes equal, and one vertical longer or shorter than these, all at right angles. The common forms are: *Prisms*, faces intersecting one or both horizontal axes, and parallel to the vertical; *Pyramids*, faces intersecting the vertical and one or both horizontal axes; *Basal pinacoids*, faces intersecting the vertical and parallel to the horizontal axes.

3. *Hexagonal* has three horizontal axes equal and making angles of 60° with each other, and one vertical, longer or shorter than these, and at right angles to them. The common forms are: *Prisms*, faces intersecting two or three of the horizontal axes and parallel to the vertical; *Pyramids*, faces intersecting the vertical and two or three of the horizontal; *Basal pinacoids*, faces intersecting the vertical and parallel to the horizontal; *Rhombohedrons*, solids of six oblique rhombic faces; *Scaleno-hedrons*, solids of twelve scalene-triangular faces.
4. *Orthorhombic* has three unequal axes all at right angles: a short forward-and-back horizontal axis, the brachyaxis; a long right-and-left horizontal axis, the macroaxis; and a vertical axis. The common forms are: *Prisms*, faces intersecting the horizontal axes and parallel to the vertical; *Pyramids*, faces intersecting the three axes; *Macropinacoids*, faces intersecting the brachyaxis and parallel to the other two; *Brachypinacoids*, faces intersecting the macroaxis and parallel to the other two; *Macrodomes*, faces intersecting the brachyaxis and the vertical and parallel to the macroaxis; *Brachydomes*, faces intersecting the macroaxis and the vertical and parallel to the brachyaxis; *Basal pinacoids*, faces intersecting the vertical and parallel to the horizontal axes.
5. *Monoclinic* has three unequal axes: a forward-and-back inclined axis, the clinoaxis; a right-and-left horizontal axis, the orthoaxis; and a vertical axis. The common forms are: *Prisms*, faces intersecting the two lateral axes and parallel to the vertical; *Pyramids*, faces intersecting all three axes; *Ortho-*

pinacoids, faces intersecting the clinoaxis and parallel to the other two; *Clinopinacoids*, faces intersecting the orthoaxis and parallel to the other two; *Orthodomes*, faces intersecting the clinoaxis and the vertical and parallel to the orthoaxis; *Clinodomes*, faces intersecting the orthoaxis and the vertical and parallel to the clinoaxis; *Basal pinacoids*, faces intersecting the vertical and parallel to the other two.

6. *Triclinic* has three unequal axes, all oblique to each other. The common forms are the same as in the orthorhombic system, namely, *Prisms*; *Pyramids*; *Macropinacoids*; *Brachypinacoids*; *Macrodomes*; *Brachydomes*; *Basal pinacoids*.

Twinning.—Some crystals instead of being simple individuals are made up of two crystals, not in parallel position, but united along a plane common to both, and such crystals are said to be twinned. Twinning is usually indicated by reentrant angles between the faces.

Cleavage.—The property which a mineral has of splitting or breaking along certain definite directions is called cleavage. The cleavage is always parallel to a possible crystal plane, and the kind of cleavage is designated by the name of the plane to which it corresponds in direction. The common kinds of cleavage for each system are:

Isometric, cubic, octahedral, and dodecahedral.

Tetragonal, basal and prismatic.

Hexagonal, basal, prismatic, and rhombohedral.

Orthorhombic, basal; prismatic; macro- or brachypinacoidal.

Monoclinic, basal; prismatic; ortho- or clinopinacoidal.

Triclinic, basal and macro- or brachypinacoidal.

The direction of cleavage can usually be determined only on the

actual crystal and not on the average massive mineral specimen. Bright, smooth cleavage faces are, however, usually present on specimens of minerals which possess good cleavage, and often they are very prominent.

Fracture.—When the direction of breakage is not definite, but occurs in any way irrespective of crystal planes, the mineral fractures. The fracture may be *even*; *uneven*; *rough*; *conchoidal* = rounded, shell-like; *splintery*; these terms referring to the kind of surface.

Tenacity.—The terms used to denote the tenacity are:

Malleable, when the mineral can be flattened by hammering.

Sectile, when it can be cut with a knife but will break in pieces by hammering.

Brittle, when it will break in pieces by hammering.

Tough, when it is difficult to break by hammering.

Structure.—Most minerals do not occur as simple individual crystals in nature, but rather as aggregates of imperfectly formed crystals, or simply as crystalline masses. Some of the terms used to describe the structure of specimens are:

Massive, when the specimen has an irregular, indefinite shape.

It may be fine or coarse granular.

Crypto-crystalline, extremely fine crystalline; *impalpable* = extremely dense, compact.

Fibrous, composed of fibers. The fibers may be parallel, radiate, or divergent in any direction.

Columnar, stout fibrous, forming columns.

Capillary, hair-like fibers.

Acicular, needle-like.

Reticulated, when the fibers cross each other, forming a net-like structure.

Mammillary, large rounded surfaces.

Reniform, kidney-shaped masses.

Botryoidal, grape-like structure or small rounded surfaces.

Geodal, cavities lined with crystals.

Drusy, rough surfaces due to innumerable small imperfect crystals.

Micaceous, thin sheets or scales, like mica.

Lamellar, thin plates.

Foliated, thin leaves.

Other terms are used to describe the structures of mineral specimens, but their meaning in general is self-evident.

ANALYTICAL KEY.

	PAGE
1. STREAK DARK GRAY OR IRON-BLACK:	
Color: Dark Gray or Black.....	10-12
Metallic White to Light Metallic Gray....	12-14
Brass; Bronze; Copper-red or Brown	14-16
2. STREAK METALLIC WHITE TO LEAD- OR STEEL-GRAY: •	
Color: Metallic White or Light Metallic Gray.....	16
3. STREAK RED OR RED-BROWN:	
Color: Red or Brown.....	18
Dark Gray or Black.....	20-22
4. STREAK YELLOW OR YELLOW-BROWN:	
Color: Red.....	22
Yellow	22-24
Brown or Black.....	24-26
Green.....	26
5. STREAK BLUE OR GREEN:	
Color: Blue, Green, or Black.....	26-28
6. STREAK UNCOLORED, WHITE OR LIGHT GRAY:	
Color: Yellow or Brown.....	28-38
Pink, Red, or Red-violet	38-44
Blue or Blue-violet.....	44-46
Green	46-54
Black.....	54-56
White, Gray, or Colorless.....	56-68

STREAK DARK GRAY

	Name.	Composition.	Color.	Streak.	Luster.	H.
COLOR DARK GRAY OR BLACK.	GRAPHITE	C	Dark steel-gray Iron-black	Black Dark silver-gray	Metallic Dull	1.2
	MOLYBDENITE	MoS ₂	Bluish lead-gray	Lead-gray Sometimes greenish	Metallic	1 1.5
	PYROLUSITE	MnO ₂	Black Blackish gray	Dull black	Metallic Dull	2
	STIBNITE	Sb ₂ S ₃	Dark lead-gray	Dark lead-gray Black	Metallic	2
	JAMESONITE	Pb ₂ Sb ₂ S ₅	Dark lead-gray	Grayish black	Metallic	2 3
	ARGENTITE	Ag ₂ S	Dark lead-gray Black	Dark lead-gray	Metallic	2 2.5
	STEPHLENITE	Ag ₅ SbS ₄	Iron-black	Iron-black	Metallic	2 2.5
	GALENITE	PbS	Dark lead-gray	Grayish black Dark lead-gray	Metallic	2.5 3
	CHALCOCITE	Cu ₂ S	Dark lead-or steel-gray Black	Dark gray	Metallic	2.5 3
	ENARGITE	Cu ₃ As ₄	Grayish black	Grayish black	Metallic	3

System.	Cleavage or Fracture.	G.	Common Structure.	Observations.
Hex.	C, basal, perfect in crystallized masses; sectile; flexible	2.2	Foliated; scaly; massive; granular; earthy	Feels greasy; plates highly flexible; inelastic; occurs with calcite; darker than molybdenite
Hex.	C, basal, very prominent; sectile; flexible	4.7	Foliated; massive; scaly; flaky	Soft and greasy like graphite but lighter colored; usually as flakes in quartz
Orth.	C, none Brittle	4.8	Fibrous; acicular; columnar; earthy; powder	Blackens fingers; often with psilomelane; darker than stibnite
Orth.	C, brachypinacoidal, perfect and prominent Brittle; slightly sectile	4.5	Fibrous; columnar; bladed; prismatic	Prisms often bent and with long shining cleavage faces; sometimes iridescent
Orth.	C, basal, prominent Brittle	5.5 6	Acicular; fibrous; capillary	Resembles stibnite, but is heavier and has cleavage faces transverse to length
Isom.	C, not important F, hackly Slightly malleable	7.3	Octahedrons; hackly masses; arborescent; reticulated	Resembles tarnished silver; often with silver, copper, barite; cuts like lead
Orth.	C, imperfect F, uneven Very brittle	6.3	Compact; massive; crystals, short prisms	Often with other silver ores; also barite, quartz, galena
Isom.	C, cubic, perfect and prominent Sectile to brittle	7.5	Cubes; cubo-octahedrons; granular; foliated	Often with sphalerite, pyrite, tetrahedrite, cerussite, anglesite, dolomite, calcite, fluorite; heavier than stibnite and never long prismatic
Orth.	C, indistinct F, conchoidal or granular Sectile	5.7	Compact; massive; crystals with deeply striated faces	Often coated with malachite; occurs with bornite, chalcopyrite, quartz, malachite, enargite
Orth.	C, prismatic and prominent Brittle	4.4	Massive	Often with chalcocite, bornite, famatinitite

STREAK DARK GRAY

	Name.	Composition.	Color.	Streak.	Luster.	H.
COLOR DARK GRAY OR BLACK.	TETRAHEDRITE TENNANTITE	$Cu_3Sb_2S_7$ $Cu_3As_2S_7$	Dark lead- or steel- gray	Dark gray	Metallic	3 4.5
	IRON	Fe	Steel-gray Black	Black	Metallic	4 5
	PSILOMELANE	MnO, H_2O	Grayish black Dull black	Brownish black	Submetallic	5 6
	ILMENITE (Menaccanite)	$(FeTi)_2O_3$	Iron-black	Brownish black	Metallic	5.5 6
	MAGNETITE	Fe_3O_4	Iron-black	Iron-black	Metallic	5.5 6.5
	FRANKLINITE	$(Fe, Mn, Zn)_2O_4$	Iron-black	Brownish black	Metallic	5.5 6.5
	COLUMBITE	$(Fe, Mn)(Nb, Ta)_2O_6$	Pitch- black	Grayish black	Submetallic Vitreous	6
COLOR METALLIC WHITE TO LIGHT GRAY.	STIBNITE	Sb_2S_3	Light lead- gray	Dark lead- gray Black	Metallic	2
	GALENITE	PbS	Lead-gray	Dark lead- gray Black	Metallic	2.5 3
	ANTIMONY	Sb	Light steel- gray Tin-white	Lead-gray	Metallic	3 3.5
	ARSENIC	As	Light steel- gray	Dark gray	Metallic	3.5

System.	Cleavage or Fracture.	G.	Common Structure.	Observations.
Isom	C, none F, granular Brittle	4.4 5.1	Compact; massive; tetrahedral crystals	Often in quartz with galena, chalcopyrite, sphalerite; sometimes with very brilliant luster
Isom.	C, not important F, hackly Malleable	7.3 7.8	Massive	Meteorites often have pitted and fused surfaces
None	C, none F, conchoidal and prominent Tough to brittle	3.7 4.7	Impalpable; massive; stalactitic; botryoidal; rounded masses	Often with limonite, pyrolusite, manganite
Hex	C, none F, conchoidal Brittle	4.5 5	Grains and pebbles; black sand; platy; massive granular.	Slightly magnetic to non-magnetic
Isom.	C, not prominent F, uneven Brittle	5.2	Octahedrons; massive granular to compact; sand	Strongly magnetic; often with quartz, feldspar, hornblende, chlorite; crystals usually very perfect
Isom.	C, none F, uneven Very brittle	5.2	Octahedrons, usually rounded; granular; massive	Usually with zincite, willemite, rhodonite, and calcite; magnetic, but not strongly like magnetite
Orth	C, not important F, uneven Brittle	5.3 7.3	Crystals, usually in parallel groups	Occurs in granite, often with albite, tourmaline, beryl
Orth.	C, brachypinacoidal very prominent Brittle; slightly sectile	4.5	Prismatic; fibrous; columnar; bladed	Often in quartz with galenite, sphalerite, tetrahedrite
Isom.	C, cubic, perfect and prominent Sectile to brittle	7.5	Cubes; cubo-octahedrons; granular; foliated; massive	Much heavier than stibnite and never long prismatic
Hex.	C, basal, prominent Brittle	6.7	Massive; lamellar	Often with stibnite; usually coated with earthy white oxide of antimony
Hex.	C, basal, not usually prominent F, granular Brittle	6	Rounded, reniform masses; granular	Usually tarnished dull black on surface

	Name.	Composition.	Color.	Streak.	Luster.	H.
COLOR METALLIC WHITE TO LIGHT GRAY.	ARSENOPYRITE	FeAsS	Light steel-gray Tin-white, often with brassy or reddish tinge	Grayish black	Metallic	5.5 6
	SMALTITE CHLOANTHITE	CoAs ₂ NiAs ₂	Tin-white Light steel- gray	Grayish black	Metallic	5.5 6
	COBALTITE	CoAsS	Silver- white with usually copper- red tinge	Grayish black	Metallic	5.5
	MARCASITE	FeS ₂	Brassy steel-gray Pale brass- yellow	Greenish black Brownish black	Metallic	6 6.5
COLOR BRASS, BRONZE, COPPER-RED, OR BROWN.	BORNITE	Cu ₃ FeS ₄	Copper- brown Horseflesh- brown	Grayish black	Metallic	3
	ENARGITE FAMATINITE	Cu ₃ AsS ₄ Cu ₃ SbS ₄	Reddish brown Bronze- brown	Grayish black	Metallic	3
	MILLERITE	NiS	Brass-yel- low	Greenish black	Metallic	3 3.5
	CHALCOPYRITE	CuFeS ₂	Deep brass- yellow	Greenish black	Metallic	3.5 4
	PYRRHOTITE	Fe ₇ S ₈ to Fe ₁₁ S ₁₂	Bronze- yellow Bronze- brown	Grayish black	Metallic	3.5 4.5

superior le pedo
 subnate ch.;, 1st pyrite, wulfenite,
 waverite,

OR IRON-BLACK.


System.	Cleavage or Fracture.	G.	Common Structure.	Observations.
Orth.	C, not prominent F, uneven Very brittle	6	Compact massive; pointed pyramids with horizontally striated faces	Often with sphalerite, galena, quartz; sometimes with gold; whiter than pyrite or marcasite; more common than smaltite
Isom.	C, not important F, granular Brittle	6.5	Compact; massive; reticulated	Often with copper-red niccolite, erythrite
Isom.	C, cubic, not prom- inent Brittle	6.3	Crystals commonly; cubes; pyritohedrons; massive	Often with smaltite, nico- colite; crystals usually tarnished to pale copper- red color
Orth.	C, not prominent F, uneven Brittle	4.9	Stalactitic with rough surfaces; cox- comb; radiate; col- umnar; low pyra- mids; massive	Never in cubes or pyrito- hedrons, and different in form from pyrite
Isom.	C, not important F, uneven Brittle	4.9 5.4	Compact; massive	Usually tarnished to pea- cock colors; occurs with quartz, chalcocite, chalco- pyrite
Orth.	C, prismatic and prominent Brittle	4.4	Massive	Often with chalcocite, bornite
Hex.	C, perfect and prom- inent in crystals Brittle	5.3 5.6	Acicular; capillary; hair tufts; compact fibrous layers	Always needle-like or fibrous; often in cavities in chert or red hematite, or coating pyrrhotite
Tetrag.	C, not important F, uneven to con- choidal Brittle	4.2	Massive; tetrahe- dral crystals	Often with pyrite, galena, sphalerite, tetrahedrite, chalcocite, dolomite, etc.; often tarnished peacock colors
Hex.	C, not important F, uneven Brittle	4.6	Massive; granular; occasional crystals	Usually slightly mag- netic; surface often tar- nished dark bronze-brown

STREAK DARK GRAY

BRASS, BRONZE, COP- PER-RED, OR BROWN.	Name.	Composition.	Color.	Streak.	Luster.	H.
	NICCOLITE	NiAs	Pale cop- per-red	Brownish black	Metallic	5 5.5
	PYRITE	FeS ₂	Pale brass- yellow	Greenish black Brownish black	Metallic	6 6.5
	MARCASITE	FeS ₂	Pale brass- yellow	Greenish black Brownish black	Metallic	6 6.5

STREAK METALLIC WHITE

COLOR METALLIC WHITE OR LIGHT METALLIC GRAY.	MERCURY	Hg	Tin-white		Metallic	
	MOLYBDENITE	MoS ₂	Bluish lead-gray	Lead-gray with often greenish tinge	Metallic	1 1.5
	SYLVANITE CALAVERITE	(AuAg)Te ₂	Silver- or tin-white; often with brassy tinge	Silver- white	Metallic	1.5 2
	BISMUTH	Bi	Reddish white to light cop- per-red	Silver- white Lead-gray	Metallic	2 2.5
	SILVER	Ag	Silver- white	Silver- white	Metallic	2.5
	ANTIMONY	Sb	Tin-white Silver- white	Silver- white	Metallic	3 3.5
	ARSENIC	As	Tin-white Light lead- or steel-gray	Tin-white	Metallic	3.5
	PLATINUM PLATINIRIDIUM	Pt PtIr	Tin-white Light steel- gray	Light steel- gray	Metallic	4 4.5



OR IRON-BLACK.

System.	Cleavage or Fracture.	G.	Common Structure.	Observations.
Hex.	C, none F, uneven Brittle	7.5	Compact; impalpable massive	Usually with smaltite
Isom.	C, indistinct F, uneven Brittle	5	Cubes; pyritohedrons; octahedrons; massive; granular	Very common; associated with all sulphides and in all rocks
Orth.	C, not important F, uneven Brittle	4.9	Coxcomb and curved dome shapes; stalactitic with rough faces	Distinguished from pyrite by form generally

TO LIGHT LEAD-OR STEEL-GRAY.

		13.6	Liquid globules	Occurs as small globules on cinnabar
Hex.	C, basal, perfect and prominent Sectile	4.7	Foliated masses; scales; flakes	Soft and greasy, like graphite; highly flexible; often with quartz
Mono.	C, clinopinacoidal, perfect, prominent F, coarse granular Brittle	9.9 8.3	Massive; crystals with deeply striated faces	Often in gray phonolite rock with purple fluorite; also in schist
Hex.	C, basal, perfect and prominent Brittle	9.7	Reticulated; embedded lenticular crystals; massive	Often as lenticular crystals or grains in quartz
Isom.	C, none F, hackly Malleable	10.1 11.1	Wires; arborescent; massive; filiform	Usually tarnished on surface to brown or black; often with barite, calcite, other silver ores
Hex.	C, basal, prominent Brittle	6.7	Massive; lamellar	Often with stibnite; usually coated with whitish oxide of antimony
Hex.	C, basal, not usually prominent F, granular Brittle	5.6	Rounded reniform masses; granular	Usually tarnished dull black on surface
Isom.	C, none F, hackly Malleable	14 19	Nuggets; grains	In gold-bearing sands

STREAK RED

COLOR RED OR BROWN.	Name.	Composition.	Color.	Streak.	Luster.	H.
	BAUXITE	$\text{Al}_2\text{O}_3 + 2\text{H}_2\text{O}$	Brown	Reddish brown	Earthy Dull	1 2
	HEMATITE	Fe_2O_3	Brownish red Cherry-red	Dark red Cherry-red	Earthy Dull	1 4
	ERYTHRITE	$\text{Co}_3\text{As}_2\text{O}_8 + 8\text{H}_2\text{O}$	Peach-red Crimson	Pale red	Earthy Vitreous	1 2.5
	WAD	$\text{MnO}, \text{H}_2\text{O}$	Dark brown	Dark reddish brown	Earthy Dull	1 3
	CINNABAR	HgS	Scarlet red Vermillion Dark red	Scarlet Vermillion	Adamantine	2 2.5
	PROUSTITE	Ag_3AsS_3	Scarlet Vermillion	Scarlet	Adamantine	2 2.5
	COPPER	Cu	Copper-red	Copper-red	Metallic	2.5 3
	SPHALERITE	ZnS	Dark brown	Reddish brown	Resinous Vitreous	3.5 4
	CUPRITE	Cu_2O	Dark red	Cochineal-red Brick-red Crimson-red	Adamantine Earthy	3.5 4
	HEMATITE	Fe_2O_3	Dark brownish red	Brownish red	Submetallic	4.5
TURGITE	$2\text{Fe}_2\text{O}_3, \text{H}_2\text{O}$	Brown	Reddish brown	Submetallic	5 6	

OR RED-BROWN.

Red hematite
red color

System	Cleavage or Fracture.	G.	Common Structure.	Observations.
	C, none F, earthy	2.5	Claylike masses with small rounded concretions; pisolitic	Clay odor; distinguished from clay by pisolitic structure
	C, none F, earthy	5	Earthy masses; oolitic; powder	Red ochre; often red clay
Mono.	C, not important F, usually earthy	2.9	Earthy; crusts; rarely in crystals	Occurs with cobalt and nickel ores as coatings and crusts
Amorph.	C, none F, earthy Brittle	4	Earthy masses; powder concretions	Often with psilomelane, limonite, malachite, azurite
Hex.	C, prismatic, not important F, uneven Sectile	8 8.2	Granular; crystals; powder; massive; compact	Occurs with marcasite, chalcedony, quartz, sulphur; very heavy; often mixed with siliceous rock and apparently hard
Hex.	C, rhombohedral, not prominent Brittle	5.6	Crystals; red bands or streaks in rock	Light ruby silver ore; often with gray pyrrargyrite
Isom.	C, none F, hackly Malleable	8.8	Hackly masses; sheets; wires, arborescent forms	Usually tarnished black on surface; often with calcite, cuprite, malachite
Isom.	C, dodecahedral, perfect and prominent Brittle	4	Massive; crystals	Often with galena, pyrite, arsenopyrite, etc.
Isom.	C, poor F, uneven Brittle	5.9	Massive; compact; crystals; octahedrons; cubes	Usually with malachite, copper
Hex.	C, none F, uneven; splintery Brittle	5	Massive; reniform, mamillary; botryoidal; splintery; oolitic	Massive red hematite
	C, none F, splintery Brittle	4.2 4.4	Compact; fibrous; massive; botryoidal; earthy	Resembles limonite; distinguished by streak; fibers often with satin-like luster

STREAK RED

Name.	Composition.	Color.	Streak.	Luster.	H.
PYRARGYRITE	Ag_3SbS_3	Dark steel-gray	Purple-red Cherry-red	Metallic	2.5
TETRAHEDRITE	$\text{Cu}_9\text{Sb}_2\text{S}_7$	Dark lead or steel gray	Cherry-red Dark red brown	Metallic	3 4.5
HEMATITE	Fe_2O_3	Dark steel- gray Iron-black	Brownish red	Metallic Brilliant	2.5 4
SPHALERITE	ZnS	Brownish black	Dark brown	Resinous Submetallic	3.5 4
MANGANITE	$\text{Mn}_2\text{O}_3 + \text{H}_2\text{O}$	Iron-black Dark steel- gray	Dark red- dish brown	Metallic	4
WOLFRAMITE	$(\text{Fe}, \text{Mn})\text{WO}_4$	Dark gray- ish or brownish black	Dark red- dish brown	Submetallic Metallic	5 5.5
CHROMITE	FeCr_2O_4	Black Brownish black	Grayish brown	Submetallic to pitch-like	5.5
PSILOMELANE	$\text{MnO}, \text{H}_2\text{O}$	Dull black	Very dark brown	Submetallic Dull	5 6
HEMATITE	Fe_2O_3	Iron-black Dark steel- gray	Cherry-red Brownish red Red-brown	Metallic	5.5 6.5
ILMENITE	$(\text{Fe}, \text{Ti})_2\text{O}_3$	Iron-black	Very dark brown	Metallic	5.5 6
FRANKLINITE	$(\text{Fe}, \text{Mn}, \text{Zn})_3\text{O}_4$	Iron-black	Dark red- dish brown Blackish brown	Metallic	5.5 6.5

COLOR DARK GRAY OR BLACK.

OR RED-BROWN.

System.	Cleavage or Fracture.	G.	Common Structure.	Observations.
Hex.	C, not important F, uneven Brittle	5.8	Massive; granular; bands and streaks in rock	Usually as gray bands mixed with red, in quartz rock; dark ruby silver ore
Isom.	C, none F, uneven Brittle	4.4 5.1	Massive; tetrahe- dral crystals	Often has brilliant luster with brassy tinge; in quartz with sulphides
Hex.	C, micaceous Brittle	4.9 5.3	Foliated; platy; micaceous	Specular hematite; very bright sparkling plates or scales
Isom.	C, dodecahedra l, prominent Brittle	4	Massive	Often with galenite, py- rite, chalcopyrite, tetra- hedrite
Orth.	C, brachypinacoid- al, prominent Brittle	4.4	Prisms; columnar; acicular	Prisms often in bunches with prism faces deeply striated vertically; occurs with pyrolusite
Mono.	C, clinopinacoidal, perfect and prominent Brittle	7.5	Thick tabular crys- tals; massive; com- pact	Often with cassiterite, quartz, fluorite
Isom.	C, none F, uneven Brittle	4.3 4.6	Massive; granular	Sometimes coated with green, garnet; often with serpentine
	C, none F, conchoidal Tough to brittle	3.7 4.7	Impalpable; mas- sive; stalactitic; bot- tryoidal, round masses	Often with powdery pyrolusite
Hex.	C, none F, uneven Brittle	4.9 5.3	Massive; granular; foliated; crystals; scales; micaceous	Crystals often have an iridescent tarnish; fine scaly specular variety seems soft
Hex.	C, none F, conchoidal Brittle	4.5 5	Rounded pebbles; sand; plates; mas- sive	Sometimes slightly mag- netic
Isom.	C, none F, uneven Very brittle	5 5.2	Rounded crystals; Octahedrons; granu- lar masses	Usually with zincite, willemite, calcite; mag- netic but not so strongly as magnetite

STREAK RED

BLACK.	Name.	Composition.	Color.	Streak.	Luster.	H.
	COLUMBITE	(Fe,Mn)(Nb,Ta) ₂ O ₆	Pitch-black	Dark brown	Submetallic Vitreous	6
CASSITERITE	SnO ₂	Black	Dark brown	Submetallic to Metallic	6 7	

STREAK YELLOW

COLOR RED.	Name.	Composition.	Color.	Streak.	Luster.	H.
	REALGAR	AsS	Bright red Orange-red	Orange yellow	Adamantine Resinous Vitreous	1.5 2
ZINCITE	ZnO	Dark red Blood-red	Orange-yellow	Vitreous	4 4.5	
COLOR YELLOW.	ORPIMENT	As ₂ S ₃	Lemon-yellow	Lemon-yellow	Adamantine Resinous Pearly	1.5 2
	REALGAR	AsS	Orange-yellow	Orange-yellow	Resinous Vitreous	1.5 2
	SULPHUR	S	Sulphur-yellow Honey-yellow Straw-yellow	Pale yellow	Resinous Greasy Vitreous	1.5 2.5
	LIMONITE	2Fe ₂ O ₃ ·3H ₂ O	Yellow	Yellow Brown	Earthy Dull	2 4
	GOLD	Au	Golden yellow	Golden yellow	Metallic	2.5 3
	SPHALERITE	ZnS	Brownish yellow	Pale yellow	Resinous	3.5 4

OR RED-BROWN.

System.	Cleavage or Fracture.	G.	Common Structure.	Observations.
Orth.	C, not important F, uneven Brittle	5.3 7.3	Crystals, usually in parallel groups	Occurs in granite, often with albite, tourmaline, beryl
Tetrag.	C, imperfect F, uneven Brittle	6.8 7.1	Massive	Often in quartz - mica rock with wolframite, fluorite

OR YELLOW-BROWN.

Mono.	C, clinopinacoidal, not prominent F, conchoidal Sectile	3.5	Massive; granular; crystals	Often with orpiment; disseminated in siliceous rock and often apparently hard
Hex.	C, basal, perfect and prominent F, uneven Brittle	5.4 5.7	Massive; lamellar; granular	Occurs with franklinite, willemite, calcite
Orth.	C, brachypinacoidal, perfect and prominent Sectile; flexible	3.5	Foliated; plates massive	Usually with realgar
Mono.	C, clinopinacoidal, not prominent F, conchoidal Brittle; sectile	3.5	Crystals; massive; granular	Often with orpiment or finely mixed quartz
Orth.	C, indistinct F, conchoidal Brittle	2	Crystals; pyramids; crusts	Often with celestite, aragonite, limestone, cinnabar, gypsum
	C, none F, earthy	3.6	Earthy masses; ochre powder	Yellow ochre; often yellow clay
Isom.	C, none F, hackly Highly malleable	15.6 19.3	Scales; flakes; leaves; grains; wires; nuggets	Usually in quartz, conglomerates, or schists; sometimes with pyrite or arsenopyrite
Isom.	C, dodecahedral; prominent F, uneven Brittle	4	Massive; cleavage masses; crystals	Usually with galena, pyrite, chalcopyrite, tetrahedrite, quartz, calcite, dolomite

STREAK YELLOW

	Name.	Composition.	Color.	Streak.	Luster.	H.
YELLOW	PYROMORPHITE	$(\text{PbCl})\text{Pb}_4\text{P}_3\text{O}_{12}$	Greenish yellow Wax-yellow	Greenish yellow	Adamantine	3.5 4
	BAUXITE	$\text{Al}_2\text{O}_3 \cdot 2\text{H}_2\text{O}$	Brown	Brownish yellow	Earthy Dull	1 3
	LIMONITE	$2\text{Fe}_2\text{O}_3 \cdot 3\text{H}_2\text{O}$	Brown	Brownish yellow Yellowish brown	Earthy Dull	2 4
COLOR BROWN OR BLACK	SPHALERITE	ZnS	Brown Brownish black	Brownish yellow	Resinous	3.5 4
	SIDERITE	FeCO_3	Pale brown Grayish brown Dark brown	Pale yellow Yellowish brown	Vitreous	3.5 4
	GOETHITE	$\text{Fe}_2\text{O}_3 \cdot \text{H}_2\text{O}$	Yellowish brown	Yellowish brown Brownish yellow	Submetallic	5 5.5
	LIMONITE	$2\text{Fe}_2\text{O}_3 \cdot 3\text{H}_2\text{O}$	Yellowish brown Dark brown	Brownish yellow Yellow-brown	Submetallic	5 5.5
	CHROMITE	FeCr_2O_4	Black	Grayish brown	Submetallic Pitchlike	5.5
	BROOKITE	TiO_2	Dark brownish black	Pale yellow Grayish brown	Submetallic Metallic	5.5 6
	RUTILE	TiO_2	Reddish brown Black	Pale yellowish brown	Adamantine Metallic	6 6.5

System.	Cleavage or Fracture.	G.	Common Structure.	Observations.
Hex.	C, not prominent F, uneven Brittle	6.5 7.1	Small hexagonal prisms; massive	Often with galena, cerussite, anglesite, mimetite
	C, none F, earthy	2.5	Clay-like masses; pisolitic	Clay odor; distinguished from kaolinite (clay) by pisolitic structure
	C, none F, earthy	3.6	Massive; earthy	Brown ochre or brown clay (kaolinite)
Isom.	C, dodecahedral, prominent F, uneven Brittle	4	Massive; cleavage masses; crystals	Common color; occurs with galena in chert; also with many sulphides
Hex.	C, rhombohedral, perfect and prominent Brittle	3.8	Rhombohedrons; cleavage masses; crystals with curved faces	Often with cryolite, quartz, hematite, fluorite
Orth.	C, brachypinacoidal, prominent F, uneven Brittle	4 4.4	Acicular; stalactitic; radiate; fibrous	Often in cavities in limonite or hematite; distinguished from limonite by crystals and cleavage
	C, none F, uneven Brittle	3.6 4	Compact; massive; stalactitic; botryoidal; columnar	Often in cubes as an alteration from pyrite; very common iron oxide; botryoidal masses often have black varnish-like surfaces
Isom.	C, none F, uneven Brittle	4.3 4.5	Massive	Often coated with green garnets; often with serpentine
Orth.	C, not important F, uneven Brittle	3.8 4	Square pyramids; hexagonal shaped pyramids	Always in crystals; faces deeply striated; not twinned like rutile
Tetrag.	C, not important F, uneven Brittle	4.2	Twinned crystals; long acicular crystals	Faces deeply striated; knee-shaped twins; often in quartz

STREAK YELLOW

	Name.	Composition.	Color.	Streak.	Luster.	H.
BR. OR BL.	CASSITERITE	SnO_2	Black	Pale yellow	Submetallic	6 7
			Reddish brown Yellowish brown Pale yellow			
COLOR GREEN.	PYROMORPHITE	$(\text{Pb}, \text{Cl})\text{Pb}_4(\text{PO}_4)_3$	Yellowish green Grass-green	Pale greenish yellow	Adamantine Vitreous	3.5 4
	EPIDOTE	$\text{HCa}_2(\text{Al}, \text{Fe})_2\text{Si}_2\text{O}_{13}$	Yellowish green Deep green Oil-green	Pale yellow	Vitreous	6 7

STREAK BLUE

COLOR BLUE, GREEN, OR BLACK.	ANNABERGITE	$\text{Ni}_3\text{As}_2\text{O}_8 + 8\text{H}_2\text{O}$	Apple-green	Pale green	Vitreous Earthy	1
	VIVIANITE	$\text{Fe}_3\text{P}_2\text{O}_8 + 8\text{H}_2\text{O}$	Dark blue-green Indigo-blue	Indigo-blue	Vitreous Earthy	1.5 2
	CHLORITE PROCHLORITE CLINOCLORE	$\text{Mg}_3\text{FeAl}_2\text{Si}_2\text{O}_{11} + 4\text{H}_2\text{O}$	Dark green	Grayish green	Vitreous Pearly	1.5 2.5
	LINARITE	$\text{CuPbSO}_4 \cdot \text{H}_2\text{O}$	Deep azure-blue	Smalt-blue Pale blue	Adamantine Vitreous	2.5
	CHRYSOCOLLA	$\text{CuSiO}_3 + 2\text{H}_2\text{O}$	Bluish green Greenish blue	Pale green Pale blue	Vitreous Greasy Earthy	2 4
	AZURITE	$2\text{CuCO}_3 \cdot \text{Cu}(\text{OH})_2$ <i>Co y. 10.</i>	Azure-blue	Smalt-blue	Vitreous Velvety	3.5 4
	MALACHITE	$\text{CuCO}_3 \cdot \text{Cu}(\text{OH})_2$	Bright green Emerald-green Dark green	Emerald-green	Vitreous Silky Velvety Dull	3.5 4

OR YELLOW-BROWN.

System.	Cleavage or Fracture.	G.	Common Structure.	Observations.
Tetrag.	C, not important F, uneven Brittle	6.8 7.1	Pebbles with concentric structure; crystals; massive	Stream tin; wood tin
Hex.	C, not important F, uneven Brittle	6.5 7.1	Small hexagonal prisms; drusy crusts; massive	Very heavy; usually with ores of lead
Mono.	C, basal, perfect and usually prominent	3.2 3.5	Prismatic; columnar; reticulated masses	Often in quartz and schists; also with calcite

OR GREEN.

Mono.	C, none F, earthy		Fine capillary coatings	Occurs with erythrite as coatings and crusts on cobalt and nickel ores
Mono.	C, clinopinacoidal, perfect in crystals Brittle	2.6	Slender prismatic; acicular; earthy	Often as crystals in pyrrhotite; as earthy round masses in clay
Mono.	C, basal, perfect and prominent Tough to brittle	2.8	Micaceous scaly flakes; compact scaly masses	Highly flexible but not elastic, like mica; often altered from biotite
Mono.	C, orthopinacoidal, usually prominent Brittle	5.4	Columnar; fibrous; long prisms sometimes radiate	Usually with galenite; heavier than azurite, and shows cleavage faces
Amorph.	C, none F, uneven Brittle	2 2.3	Massive; stains; earthy	Usually with copper ores; darker and glassier bluish green than malachite, and never fibrous
Mono.	C, not prominent Brittle	3.8	Crystals; fibrous; acicular	Usually with malachite and often with limonite, wad
Mono.	C, not prominent F, uneven; splintery Brittle	4	Fibrous; banded; stalactitic; botryoidal; powder	Often with cuprite, copper, chalcocite, chalcopyrite; often as green stains in ore rocks

STREAK BLUE

COLOR BLUE, GREEN, OR BLACK.	Name.	Composition.	Color.	Streak.	Luster.	H.
	ALABANDITE	MnS	Black	Dark grayish green	Submetallic	3.5 4
	LAZURITE LAPIS LAZULI	NaAlSi ₃ O ₈	Azure-blue Ultramarine blue	Pale blue	Vitreous Greasy	5 5.5
	AUGITE	MgCa ₂ FeSi ₆ O ₁₂	Greenish black Blackish green	Pale grayish green	Vitreous	5 6
	HORNBLLENDE	Mg ₃ Ca ₂ FeSi ₆ O ₁₈	Greenish black Blackish green	Pale grayish green	Vitreous Silky	5 6
	GLAUCOPHANE	Silicate of Na, Al, Mg, Fe	Lavender-blue Blackish blue	Grayish blue	Vitreous Pearly	6 6.5

STREAK UNCOLORED, WHITE,

COLOR YELLOW OR BROWN.	Name.	Composition.	Color.	Luster.	H.
	CERARGYRITE	AgCl	Dark gray Dark brown	Waxy Adamantine	1 1.5
	BAUXITE	Al ₂ O ₃ +3H ₂ O	Yellow to brown	Earthy Dull	1.5 3
	KAOLINITE (Clay)	H ₄ Al ₂ Si ₂ O ₉	Yellow to brown	Earthy Dull	1.5 2.5
	GYPSUM	CaSO ₄ +2H ₂ O	Yellow to brown	Vitreous Silky Dull	1.5 2
	SULPHUR	S	Sulphur-yellow Honey-yellow Brown	Vitreous Greasy	1.5 2

OR GREEN.

29

System.	Cleavage or Fracture.	G.	Common Structure.	Observations.
Isom.	C, not prominent F, uneven Brittle	4	Massive	Often with rhodochro- site, pyrite, argentite, ga- lena; surface usually tar- nished brown
Isom.	C, not important Brittle	2.4	Massive; dissemi- nated in rock	Often with calcite and pyrite
Mono.	C, prismatic, not usually promi- nent Cleavage angle 87° Brittle	3.2 3.6	Almost square prisms; massive	Distinguished from horn- blende by cleavage angle; also more often in crystals
Mono.	C, prismatic and very prominent Cleavage angle 124° Brittle	2.9 3.4	Massive; crystals rare	Usually with bright cleav- age faces having a fibrous appearance
Mono.	C, prismatic, per- fect Brittle	3.1	Fibrous; columnar; reticulated	Usually forms schists; a blue hornblende

OR LIGHT GRAY.

System.	Cleavage or Fracture.	G.	Common Structure.	Observations.
Isom.	C, none Very sectile	5.5	Thin crusts; coat- ings; massive	Cuts like wax; often with silver ores
	C, none F, earthy	2.5	Compact earthy; pisolitic	Clay odor; distinguished from kaolinite (clay) by pea-shaped structure
Mono.	C, none F, earthy	2.6	Compact earthy; soapy; friable	Clay odor; massive clay; occasionally soapy feel
Mono.	C, pinacoidal, per- fect and promi- nent Brittle	2.3	Fibrous; columnar; granular; compact massive	Ferruginous gypsum
Orth.	C, not important F, uneven Brittle	2	Crystals; pyramids; crusts.	Often with limestone, celestite, aragonite, cinna- bar

STREAK UNCOLORED, WHITE,

Name.	Composition.	Color.	Luster.	H.
CHLORITE PROCHLORITE CLINOCHLORE PENNINITE, etc.	$H_2(Mg, Fe)_4Al_2Si_2O_{13}$	Dark yellowish brown Greenish brown	Vitreous Pearly	1.5 2.5
BIOTITE	$(HK)_2(Mg, Fe)_2(Al, Fe)_2(SiO_4)_3$	Dark brown Greenish brown	Pearly to Vitreous	2.5 3
PHLOGOPITE	$H_2KMg_3Al(SiO_4)_3$	Light brown Cinnamon-brown	Pearly Vitreous	2.5 3
HALITE	NaCl	Light yellow or brown	Vitreous	2.5
CRYOLITE	Na_3AlF_6	Grayish brown	Vitreous Icy	2.5 3
CALCITE	$CaCO_3$	Honey-yellow Light to dark brown	Vitreous	3
BARITE	$BaSO_4$	Lemon-yellow Yellowish brown	Vitreous	2.5 3.5
SERPENTINE	$(H_2(MgOH)Mg_2(SiO_4)_2$	Greenish brown Yellowish brown	Greasy Vitreous	2.5 4
CERUSSITE	$PbCO_3$	Grayish brown Yellowish brown	Adaman- tine Earthy	3 3.5
WULFENITE	$PbMoO_4$	Lemon-yellow Orange-yellow	Greasy Adaman- tine	3
MIMETITE	$(PbCl)Pb_4(AsO_4)_3$	Brownish yellow Yellow-brown	Adaman- tine Greasy	3.5
PYROMORPHITE	$(PbCl)Pb_4(PO_4)_3$	Greenish yellow Yellowish brown	Greasy Adaman- tine	3.5 4

COLOR YELLOW OR BROWN.

System.	Cleavage or Fracture.	G.	Common Structure.	Observations.
Mono.	C, basal, perfect and prominent Tough	2.9	Micaceous; scales; flakes; compact scaly masses	Highly flexible but not elastic, like the micas; often an altered biotite
Mono.	C, basal, perfect and very prominent Tough	2.7 3.1	Mica plates; scales; flakes	Dark color even in thinnest plates; flexible and elastic; black mica
Mono.	C, basal, perfect and prominent Tough	2.8	Micaceous; scales; flakes; sheets	Lighter brown than biotite; almost colorless in thin sections; brown mica
Isom.	C, cubic, perfect and prominent Brittle	2.1 2.6	Cubes; massive; granular	Tastes salt
Mono.	C, basal, prominent, and pinacoidal not so good	3	Massive	Three cleavages almost at right angles, making cubes; usually with siderite
Hex.	C, rhombohedral, very prominent Brittle	2.7	Rhombohedrons; scalenohedrons; granular; massive	Rhombohedral cleavage very characteristic; very common mineral
Orth.	C, basal and prismatic, perfect and prominent	4.3 4.6	Massive; platy crystals	Heavy, vitreous mineral; often with galena
Mono.	C, not important Brittle	2.6	Massive; compact	Very smooth feel, almost greasy
Orth.	C, not prominent F, conchoidal Very brittle	6.5	Massive; crusts	Usually with galenite or anglesite; very heavy
Tetrag.	C, not prominent Brittle	6.7 7	Square plates; thin plates and tables	Often with vanadinite or galenite
Hex.	C, not prominent Brittle	7 7.2	Rounded aggregates of plates; small crystals	Often with pyromorphite, galena
Hex.	C, none Brittle	6.5 7.1	Short hexagonal prisms; columnar masses	Often with galena, cerussite, anglesite; crystal faces deeply striated vertically

STREAK UNCOLORED, WHITE,

COLOR YELLOW OR BROWN.	Name.	Composition.	Color.	Luster.	H.
	ARAGONITE	CaCO_3	Honey-yellow Yellow-brown	Vitreous Glassy	3.5 4
	STILBITE	$(\text{Na}_2\text{Ca})\text{Al}_2\text{Si}_6\text{O}_{18}\cdot 6\text{H}_2\text{O}$	Yellowish brown Light brown	Vitreous Silky	3.5 4
	DOLOMITE	$(\text{CaMg})\text{CO}_3$	Yellowish brown Grayish brown	Vitreous	3.5 4
	SIDERITE	FeCO_3	Grayish brown Dark brown	Vitreous Pearly	3.5 4
	SPHALERITE	ZnS	Honey-yellow Yellowish brown Reddish brown	Resinous	3.5 4
	MAGNESITE	MgCO_3	Grayish brown	Vitreous Dull	3.5 4.5
	FLUORITE	CaF_2	Lemon-yellow Pale yellow Yellowish brown	Vitreous Glassy	4
	SCHEELITE	CaWO_4	Yellowish brown Grayish brown	Greasy Adamantine	4.5 5
	CALAMINE	$\text{H}_2\text{Zn}_2\text{SiO}_5$	Pale brown	Vitreous	4.5 5
	SMITHSONITE	ZnCO_3	Yellowish brown	Vitreous	5
APATITE	$(\text{CaF})\text{Ca}_4(\text{PO}_4)_3$	Brown Greenish brown	Vitreous Greasy	5	

System.	Cleavage or Fracture.	G.	Common Structure.	Observations.
Orth.	C, not prominent Brittle	2.9	Stalactitic; banded; massive	Differs from calcite in not having prominent cleavage
Mono.	C, clinopinacoidal Brittle	2 2.2	Columnar; sheaf- like; fibrous	Often in cavities in lava rocks, with chabazite, heu- landite, analcite
Hex.	C, rhombohe d r a l , not prominent	2.9	Massive	Harder than calcite *
Hex.	C, rhombohe d r a l , perfect and very prominent	3.8	Rhomb o h e d r o n s with curved faces; s a d d l e - s h a p e d c r y s - t a l s ; m a s s i v e	Often in cryolite; more glassy than sphalerite
Isom.	C, dodeca h e d r a l , very prominent	3.9 4.1	Massive	Resinous cleavage faces characteristic; often with galena, tetrahedrite, etc.
Hex.	C, rhombohe d r a l , prominent in crystals F, conchoidal	3.1	Massive; rhombohe- drons	Crystals usually in talc; compact impalpable mass- es more common
Isom.	C, octahedral, per- fect and promi- nent Very brittle	3.1	Cubes; granular, massive	Often with pyrite, galena, and sulphides
Tetrag.	C, not prominent Brittle	5.9 6.1	Crystals; pyramids; massive	Often with wolframite, cassiterite; very heavy
Orth.	C, prismatic, prom- inent	3.5	Drusy coatings; small crystals	Often with smithsonite on yellow earthy masses
Hex.	C, rhombohe d r a l , not prominent	4.3 4.5	Small rhombohe- dral crystals; drusy crystals; bone-like masses	Dry bone; often with the silicate calamine; also with sphalerite
Hex.	C, basal, imperfect, not prominent Brittle	3.2	Hexagonal prisms; granular	Green and brown colors often intermixed; crystals often have fused appear- ance

		Name.	Composition.	Color.	Luster.	H.
		MONAZITE	$(Ce, La, Di, Th)PO_4$	Honey-yellow Brown	Adamantine Vitreous	5 5.5
		TITANITE	$CaTiSiO_5$	Dark brown Brownish yellow	Adamantine Greasy Vitreous	5 5.5
		WILLEMITE TROOSTITE	Zn_2SiO_4	Greenish yellow Reddish brown	Vitreous	5.5
		OPAL	$SiO_2 + nH_2O$	Yellow Brown	Waxy Vitreous	5.5 6.5
COLOR YELLOW OR BROWN.	PYROXENE	ENSTATITE	$MgSiO_3$	Grayish brown Greenish brown	Vitreous Pearly	5.5
		BRONZITE	$(Mg, Fe)SiO_3$	Bronze-brown	Vitreous Bronzy	5 6
		HYPERSTHENE	$(Fe, Mg)SiO_3$	Dark brown Blackish brown	Bronze-metallic Vitreous Pearly	5 6
		AUGITE	Silicate of Ca, Mg, Al, and Fe, chiefly	Dark brown	Vitreous	5 6
AMPHIBOLE	ANTHOPHYLLITE	$(Mg, Fe)SiO_3$	Light grayish brown Brownish gray Greenish gray	Vitreous Pearly	5.5 6	
	TREMOLITE	$CaMg_3(SiO_3)_4$	Grayish brown	Vitreous Silky	5 6	
	HORNBLende	$Ca(MgFe)_2(SiO_3)_4$, with $(MgFe)_3(AlFe)Si_2O_{12}$ and $Na_2Al_2(SiO_3)_4$	Dark reddish brown	Vitreous	5 6	

OR LIGHT GRAY.

System.	Cleavage or Fracture.	G.	Common Structure.	Observations.
Mono.	C, basal, not prominent Brittle	5 5.5	Yellow sand; brown crystals	Commonly as yellow sand; crystals rare
Mono.	C, indistinct usually Brittle	3.5	Flat wedge-shaped crystals; massive	Often in syenitic rocks; often with hornblende and magnetite; masses show cleavage
Hex.	C, basal, sometimes prominent	3.9 4.1	Crystals; granular; massive	Often in calcite with zincite, franklinite
Amorph.	C, none F, conchoidal and prominent	2.2	Massive; wood-like	Softer than brown jasper; wood opal shows wood structure
Orth.	C, prismatic and brachypinacoidal, very prominent Brittle	3.2	Bladed, columnar; massive	Often softer because of alteration to serpentine; pearly cleavage faces usual
Orth.	C, prismatic and brachypinacoidal, very prominent Brittle	3.5	Reticulated masses; columnar	Bronze luster and color characteristic
Orth.	C, clinopinacoidal, very prominent Brittle	3.5	Broad cleavage; masses	More bronze brown than hornblende
Mono.	C, prismatic, not usually prominent Cleavage angle 87°	3.5	Crystals	Distinguished from hornblende by the prism being nearly square
Orth.	C, prismatic and prominent	3.2	Lamellar; fibrous; reticulated; columnar	Often soft because of alteration; structure resembles actinolite
Mono.	C, prismatic and prominent	2.9 3.4	Bladed; columnar; prismatic; fibrous	Cleavage angle 124°; often in marble or calcite with brown tourmaline
Mono.	C, prismatic, very prominent	3.4	Crystals	Prismatic cleavage angle about 124°

COLOR YELLOW OR BROWN.	Name.	Composition.	Color.	Luster.	H.
	NEPHELITE (Elaolite)	NaAlSiO_4	Reddish brown	Greasy Vitreous	5.5 6
	ALLANITE	$(\text{CaFe}_2)(\text{Al,Fe,Ce})_2(\text{AlOH})(\text{SiO}_4)_3$	Dark brown Blackish brown	Pitchy Submetallic	5.5 6
	SILLIMANITE (Fibrolite)	Al_2SiO_5	Light grayish brown Hair-brown	Vitreous	6 7
	ZOISITE	$\text{Ca}_2\text{Al}_2(\text{AlOH})(\text{SiO}_4)_3$	Grayish brown Yellowish brown	Vitreous	6 6.5
	EPIDOTE	$\text{HCa}_2(\text{Al,Fe})_3\text{Si}_3\text{O}_{13}$	Oil brown Greenish brown Greenish yellow	Vitreous	6 7
	RUTILE	TiO_2	Reddish brown	Adamantine	6 6.5
	CASSITERITE	SnO_2	Reddish brown Yellowish brown	Adamantine Dull	6 7
	CHONDRODITE	$\text{Mg}_2[\text{Mg}(\text{F,OH})_2(\text{SiO}_4)_2]$	Reddish brown Brownish yellow	Vitreous	6 6.5
	AXINITE	$\text{AlCa}_3(\text{AlOH})(\text{BO}_3)\text{Si}_4\text{O}_{12}$	Clove-brown Yellow	Vitreous	6.5 7
	ORTHOCLASE	KAlSi_3O_8	Pale brown Flesh-brown	Vitreous Pearly	6 6.5
	QUARTZ var. Citron Smoky Ferruginous	SiO_2	Brownish yellow Hair brown Smoky brown Yellowish brown Reddish brown	Vitreous Glassy Greasy	7
CHALCEDONY var. Agate Jasper Flint	SiO_2	Brown or yellow in all shades	Waxy Vitreous	7	

System.	Cleavage or Fracture.	G.	Common Structure.	Observations.
Hex.	C, prismatic, not prominent Brittle	2.6	Massive	Usually in crystalline rock with feldspar, biotite, leucite
Mono.	C, not important F, uneven Brittle	3.5 4.2	Thin tabular crystals; seldom massive	In granitic rocks as thin brownish, pitch-like crystals and streaks
Orth.	C, brachypinacoidal, prominent Brittle	3.2	Long slender prisms; fibers; columns	Always in schist rocks; fibers often bent and reticulated
Orth.	C, brachypinacoidal, prominent Brittle	3.2	Stout columns; sometimes fibrous masses	Columnar crystals usually much broken by cross-fracture
Mono.	C, basal, perfect, sometimes prominent Brittle	3.2 3.5	Prismatic crystals; columnar; fine granular	Often as prisms in quartz
Tetrag.	C, prismatic, not important Brittle	4.2	Twinned crystals; long prisms; acicular	Crystals usually with deeply striated faces; knee-shaped twins
Tetrag.	C, not important F, uneven Brittle	6.8 7	Crystals; rounded pebbles	Pebbles of reddish, grayish, and yellowish color, with often concentric structure
Mono.	C, not prominent F, uneven, brittle	3.2	Crystals; embedded grains in rock	Often in calcite with octahedrons of spinel
Tric.	C, not prominent F, conchoidal Brittle	3.2	Thin sharp-edged crystals	Often with garnet, tourmaline, albite, quartz
Mono.	C, basal and clinopinacoidal, very prominent	2.6	Crystals; cleavage pieces; massive	Commonly with quartz, mica, hornblende; two cleavages at R. A.
Hex.	C, none F, conchoidal Very prominent Brittle	2.6	Hexagonal prisms and pyramids; prism faces usually horizontally striated; massive.	Smoky quartz common in granites with orthoclase, biotite, or hornblende; conchoidal fracture characteristic
Hex.	C, none F, conchoidal Very prominent Brittle	2.6	Colloidal masses; often banded; botryoidal; mammillary	Conchoidal fracture, very characteristic

	Name.	Composition.	Color.	Luster.	H.
COLOR YELLOW OR BROWN.	VESUVIANITE	$MgCa_2(AlOH)Al_2(SiO_4)_5$	Dark brown Greenish brown Brownish yellow Greenish yellow	Vitreous Greasy	6.5 7
	GARNET var. Grossularite Andradite Almandite Spessartite Pyrope	$Ca_3Al_2(SiO_4)_3$ $Ca_2Fe_2(SiO_4)_3$ $Fe_3Al_2(SiO_4)_3$ $Mn_3Al_2(SiO_4)_3$ $Mg_3Al_2(SiO_4)_3$	Reddish brown Yellowish brown Reddish yellow Brownish yellow	Vitreous	6.5 7.5
	TOURMALINE	$7H_2O.2Na_2O.12MgO.6B_2O_3-13Al_2O_3.24SiO_2$	Cinnamon-brown Dark brown	Vitreous Glassy	7 7.5
	STAUROLITE	$HFeAl_2Si_2O_{12}$	Dark reddish brown	Vitreous	7 7.5
	BERYL	$Be_3Al_2(SiO_6)_3$	Golden yellow Greenish yellow	Vitreous Glassy	7.5 8
	ZIRCON	$ZrSiO_4$	Slate-brown Light brown Dark brown	Vitreous Pearly Resinous	7.5 8
	TOPAZ	$Al_2(F,OH)_2SiO_4$	Honey-yellow Wine-yellow Yellowish brown	Vitreous	8
PINK, RED, OR RED-VIOLET.	GYPSUM	$CaSO_4 + 2H_2O$	Brick-red	Vitreous Silky	1.5 2
	LEPIDOLITE	$(Li,K)_2Al_2(F,OH)_2Si_3O_9$	Pale pink to Deep rose-red	Pearly	2.5 4
	VANADINITE	$(PbCl)Pb_4(VO_4)_3$	Bright red Orange-red Ruby-red	Adamantine Greasy	2.5 3
	WULFENITE	$PbMoO_4$	Orange-red	Adamantine Greasy	3
	CALCITE	$CaCO_3$	Pink Brick-red	Vitreous	3

OR LIGHT GRAY.

System.	Cleavage or Fracture.	G.	Common Structure.	Observations.
Tetrag.	C, not prominent Brittle	3.4	Square prisms with low pyramids; massive; granular	Often in white or blue calcite; prism faces generally vertically striated
Isom.	C, none F, uneven, coarse Brittle	3.1 4.3	Crystals; dodecahedrons with icosatetrahedrons; granular; massive	Often in schists and gneisses; also with calcite; usually in crystals
Hex.	C, none F, uneven Very brittle	3 3.2	Trigonal; prisms with vertically striated faces	Prisms usually much cross-fractured; often in calcite with tremolite
Orth.	C, not important F, uneven Brittle	3.7	Crystals; often twinned in crosses, or X-shaped	Usually in schists; often with cyanite; sillimanite
Hex.	C, not important F, uneven Brittle	2.6 2.8	Hexagonal prisms with base	Harder than quartz, and crystals have basal planes
Tetrag.	C, none F, conchoidal Brittle	4.7	Square prisms with pyramids; rounded grains	Often in granitic rocks; crystals always, and usually small
Orth.	C, basal, very prominent Brittle	3.4 3.6	Crystals; prisms; pyramids	Always in crystals; sometimes in cavities in rhyolite
Mono.	C, clinopinacoidal, perfect and prominent Brittle	2.3	Columnar; fibrous; massive; granular	Gypsum stained by ferric oxide
Mono.	C, basal, perfect and prominent Tough	2.9	Micaceous; flakes; scales; compact scaly masses	Usually with rose-red tourmaline, feldspar, or quartz
Hex.	C, not important Brittle	6.6 7.2	Small hexagonal prisms	Often with wulfenite or galenite
Tetrag.	C, not important Brittle	6.7 7	Square tabular crystals	Often with vanadinite
Hex.	C, rhombohedral, prominent Brittle	2.7	Rhombohedrons; stalactites; massive	Color due to stain of ferric oxide or manganese oxide

STREAK UNCOLORED, WHITE,

Name.	Composition.	Color.	Luster.	H.
HEULANDITE	$H_4CaAl_2(Si_6O_{18}) + 3H_2O$	Deep brick-red	Pearly	3.5 4
SPHALERITE	ZnS	Brownish red Yellowish red	Resinous	3.5 4
DOLOMITE	$(Ca, Mg)CO_3$	Pale pink	Vitreous	3.5 4
RHODOCHROSITE	$MnCO_3$	Rose-red	Vitreous	3.5 4.5
MARGARITE	$H_2CaAl_4Si_2O_{12}$	Pink Rose-red	Pearly Vitreous	3.5 4.5
FLUORITE	CaF_2	Violet-red Purple Pink Amethystine	Vitreous Glassy	4
CHABAZITE	$Ca_3Al_2(SiO_4)_3(Si_2O_6)_3 + 18H_2O$	Pale brick-red Flesh-red	Vitreous	4 5
APOPHYLLITE	$H_{12}Ca_2(CaOF)_2(Si_2O_7)_3$	Pale violet-red	Vitreous Pearly	4.5 5
SCAPOLITE WERNERITE	$Ca_4Al_6Si_6O_{28}$ with $Na_4Al_3ClSi_6O_{24}$	Lilac-red Violet-red Pink	Vitreous Greasy	5.5
RHODONITE	$MnSiO_3$	Rose-red Brownish red	Vitreous	5.5 6.5
OPAL	$SiO_2 + nH_2O$	Brownish red	Waxy	5.5 6.5

COLOR PINK, RED, OR RED-VIOLET.

System.	Cleavage or Fracture.	G.	Common Structure.	Observations.
Mono.	C, clinopinacoidal, prominent Brittle	2.2	Tabular plates; crystals	Often in cavities of lava rock with stilbite, chabazite, analcite
Isom.	C, dodecahedral, very prominent Brittle	3.9 4.1	Crystals; massive	Cleavage masses common; occurs with various sulphides
Hex.	C, rhombohedral, not usually prominent Brittle	2.9	Rhombohedrons; with curved faces; saddle-shaped crystals	Often with galenite, calcite quartz, chalcopyrite
Hex.	C, rhombohedral, very prominent Brittle	3.4 3.6	Rhombohedrons; massive	Often with silver ores, also quartz, galenite, pyrite
Mono.	C, basal, perfect and prominent Brittle	3	Micaceous; foliated	Often as veins in green chlorite with diaspore, corundum; not elastic like muscovite; called brittle mica
Isom.	C, octahedral, perfect and prominent Brittle	3 3.2	Cubes; massive	Often with cassiterite, wolframite, galenite, pyrite
Hex.	C, not prominent F, uneven Brittle	2	Crystals; rhombohedrons	Crystals almost cubes in shape; often with stilbite and heulandite in cavities in lava
Tetrag.	C, basal, perfect and prominent Brittle	2.4	Square prisms with base; resemble cubes	Prism faces vertically striated; basal planes have very pearly luster
Tetrag.	C, not prominent Brittle	2.8 2.8	Massive; coarse granular	Harder than fluorite
Tric.	C, prismatic, prominent Tough	3.4 3.6	Massive granular; crystals	Often in calcite with franklinite; also with silver ores
Amorph.	C, none F, conchoidal, prominent	1.9 2.3	Colloidal masses	Conchoidal fracture characteristic; softer than jasper

STREAK UNCOLORED, WHITE,

Name.	Composition.	Color.	Luster.	H.
ORTHOCLASE	$KAlSi_3O_8$	Brick-red Flesh-red	Vitreous Pearly	6 6.5
ZOISITE var. Thulite	$Ca_2Al_2(AlOH)(SiO_4)_3$	Bright rose-red	Vitreous	6 6.5
CHONDRODITE	$Mg_2[Mg(F,OH)_2](SiO_4)_2$	Dark red Brownish red	Vitreous	6 6.5
RUTILE	TiO_2	Dark red	Adamantine	6 6.5
QUARTZ var. Amethyst Rose Ferruginous	SiO_2	Amethystine Rose-red Brick-red Violet-red	Vitreous Glassy Greasy	7
CHALCEDONY var. Agate Carnelian Jasper	SiO_2	Bright red Carnelian-red Dark red Brownish red	Waxy Vitreous	7
GARNET var. Grossularite } Essonite } Andradite } Pyrope } Almandite } Spessartite }	$Ca_2Al_2(SiO_4)_3$ $Ca_2Fe_2(SiO_4)_3$ $Mg_3(Fe,Al)_2(SiO_4)_3$ $(Mg,Fe)_3(Fe,Al)_2(SiO_4)_3$ $Mn_3(Fe,Al)_2(SiO_4)_3$	Light to dark red Brownish red Cinnamon-red Rose-red	Vitreous	6.5 7.5
TOURMALINE	$H_3(Na,Li)_4Al_10B_3Si_{12}O_{33}$	Pink Rose-red	Vitreous Glassy	7 7.5
ANDALUSITE	Al_2SiO_5	Pink Pale rose	Vitreous	7 7.5
STAUROLITE	$HFeAl_2Si_2O_{13}$	Dark brownish red	Vitreous	7 7.5
SPINEL	$MgAl_2O_4$	Ruby-red	Vitreous	8

COLOR PINK, RED, OR RED-VIOLET.

OR LIGHT GRAY.

System.	Cleavage or Fracture.	G.	Common Structure.	Observations.
Mono.	C, basal and clinopinacoidal, prominent	2.6	Crystals; massive	Occurs with quartz, mica, hornblende in red granite; see feldspar
Orth.	C, brachypinacoidal, not prominent Brittle	3.3	Massive	Not common color
Mono.	C, not prominent F, uneven Brittle	3.2	Crystals; embedded grains	Occurs with spinel in crystalline limestone; often with chlorite
Tetrag.	C, not prominent F, uneven Brittle	4.2	Crystals; long slender prisms; acicular	Often as acicular crystals in quartz
Hex.	C, none F, conchoidal, prominent Brittle	2.6	Hexagonal prisms and pyramids; massive	Ferruginous quartz usually with specular hematite; rose quartz usually massive; amethyst usually in crystals
Hex.	C, none F, conchoidal, prominent Brittle to tough	2.6	Massive; cryptocrystalline; banded	Very common as Jasper; agate usually finely banded
Isom.	C, not prominent F, uneven Brittle	3.1 4.3	Crystals; granular; rounded grains; massive	Common in schists, gneisses, and crystalline limestone
Hex.	C, none F, uneven Very brittle	3 3.2	Prismatic, often radiate or divergent; long trigonal prisms	Usually in lepidolite; crystals often parti-colored red and green
Orth.	C, not prominent Brittle	3.2	Crystals; nearly square prisms; massive	Often in schists with albite, staurolite
Orth.	C, imperfect Brittle	3.7	Crystals; often twinned into crosses and X shapes	Occurs in schists with cyanite, sillimanite, andalusite, chlorite
Isom	C, imperfect Brittle	3.5 4.1	Rounded grains; small octahedrons	Resembles red garnet and ruby corundum

STREAK UNCOLORED, WHITE,

	Name.	Composition.	Color.	Luster.	H.
COLOR BLUE OR BLUE-VIOLET.	TOPAZ	$\text{Al}_2(\text{F},\text{OH})_2\text{SiO}_4$	Pink	Vitreous	8
	CORUNDUM	Al_2O_3	Ruby-red	Vitreous	9
	VIVIANITE	$\text{Fe}_3\text{P}_2\text{O}_8 + 8\text{H}_2\text{O}$	Greenish blue Indigo-blue	Vitreous Pearly Dull	1.5 2
	CHALCANTHITE	$\text{CuSO}_4 + 5\text{H}_2\text{O}$	Sky-blue Greenish blue	Vitreous	2.5
	CHRYSOCOLLA	$\text{CuSiO}_3 + 2\text{H}_2\text{O}$	Greenish blue	Greasy Vitreous Dull	2 4
	CALCITE	CaCO_3	Sky-blue	Vitreous	3
	CELESTITE	SrSO_4	Light sky-blue	Vitreous	3 3.5
	BARITE	BaSO_4	Pale greenish blue	Vitreous	2.5 3.5
	FLUORITE	CaF_2	Violet-blue Greenish blue	Vitreous Very glassy	4
	CALAMINE	$\text{H}_2\text{Zn}_2\text{SiO}_5$	Pale blue	Vitreous Silky	4.5 5
	LAZULITE	$\text{MgAl}_2\text{P}_2\text{O}_8 + \text{H}_2\text{O}$	Smalt-blue Sky-blue Azure-blue	Vitreous	5 6
	LAZURITE (Lapis Lazuli)	$\text{Na}_4(\text{AlS}_2\text{Na})\text{Al}_3(\text{SiO}_4)_3$	Deep azure-blue Berlin blue Ultramarine blue	Vitreous	5 5.5

System.	Cleavage or Fracture.	G.	Common Structure.	Observations.
Orth.	C, basal, very perfect and prominent Brittle	3.4 3.6	Crystals	Usually artificially colored; uncommon color in nature
Hex.	C, rhombohedral, not prominent Brittle to tough	3.9 4.1	Crystals; grains; massive	Often intermixed with blue when massive; very hard
Mono.	C, pinacoidal, prominent in crystals	2.6	Long prisms; fibrous; earthy	Earthy globular masses in clay or rock, common; prisms in pyrrhotite cavities
Tric.	C, not prominent F, conchoidal Brittle	2.1 2.3	Crystals; massive; stalactitic; fibrous	Taste metallic, nauseous; artificial crystals common as blue vitriol
Amorph.	C, none F, conchoidal Sectile	2 2.2	Granular; stains; incrustations; seams	Often with clay, chalcocopyrite, limonite, malachite
Hex.	C, rhombohedral, very prominent Brittle	2.7	Coarsely granular; coarse cleavage masses	Often with vesuvianite, pyroxene
Orth.	C, basal and prismatic, prominent	3.9	Massive; fibrous	Massive varieties show good cleavage and are almost colorless; heavy
Orth.	C, not prominent F, fibrous	4.3 4.6	Fibrous	Heavy fibrous mineral
Isom.	C, octahedral, very prominent Brittle	3 3.2	Cubes; compact or granular; massive	Usually violet-blue or greenish blue; often with galena, cassiterite
Orth.	C, prismatic, sometimes prominent Brittle	3.5	Drusy crystals; coatings; massive	Geodal-shaped masses with drusy surface
Mono.	C, not prominent F, uneven Brittle	3	Crystals; acute pointed pyramids	Usually as crystals in white quartzite rock
Isom.	C, not prominent Brittle	2.4	Massive	Usually intermixed with calcite and pyrite

STREAK UNCOLORED, WHITE,

	Name,	Composition.	Color.	Luster.	H.
COLOR BLUE OR BLUE-VIOLET.	SODALITE	$\text{Na}_4(\text{Al,Cl})\text{Al}_2\text{Si}_3\text{O}_{12}$	Lavender-blue Azure-blue	Vitreous Greasy	5.5 6
	OPAL	$\text{SiO}_2 + n\text{H}_2\text{O}$	Pale grayish blue Greenish blue	Waxy Vitreous	5.5 6.5
	CYANITE	Al_2SiO_5	Sky-blue Pale greenish blue	Vitreous Pearly	5 7
	TURQUOIS	$\text{AlPO}_4 \cdot \text{Al}(\text{OH})_3 + \text{H}_2\text{O}$	Greenish blue	Dull Waxy	6
	QUARTZ	SiO_2	Grayish blue Greenish blue	Vitreous Glassy	7
	CHALCEDONY	SiO_2	Grayish blue Greenish blue	Waxy Greasy	7
	CORDIERITE (Iolite)	$\text{Al}_2\text{Mg}_4(\text{AlOH})_2(\text{Si}_2\text{O}_7)_2$	Grayish blue Greenish blue Smoky blue	Vitreous Glassy	7 7.5
	BERYL	$\text{Be}_3\text{Al}_2(\text{SiO}_4)_6$	Aquamarine blue Pale blue Sky-blue	Vitreous Glassy	7.5 8
	TOPAZ	$\text{Al}_2(\text{F,OH})_2\text{SiO}_4$	Greenish blue Sky-blue	Vitreous	8
	CORUNDUM	Al_2O_3	Grayish blue Sapphire-blue	Vitreous	9
COLOR GREEN.	TALC	$\text{H}_2\text{Mg}_3(\text{SiO}_3)_4$	Pale green Deep green	Greasy	1 1.5
	VIVIANITE	$\text{Fe}_3\text{P}_2\text{O}_8 + 8\text{H}_2\text{O}$	Bluish green	Vitreous Pearly Dull	1.5 2

OR LIGHT GRAY.

System.	Cleavage or Fracture.	G.	Common Structure.	Observations.
Isom.	C, dodecahedral, not prominent Brittle	2.1 2.5	Massive; grains	Usually with nephelite, leucite and albite in syenite rock
Amorph.	C, none F, conchoidal, prominent	1.9 2.3	Massive	Color not usually homogeneous
Tric.	C, pinacoidal, prominent Tough	3.6	Bladed; reticulated	Cleavage faces usually wavy or bent and with much cross parting; often in schists with stauroilite
None.	C, none F, uneven	2.6	Irregular-shaped masses; grains; seams; impalpable	Occurs intermixed with rock in veins, seams, etc.
Hex.	C, none F, conchoidal and prominent	2.6	Crystals; massive	Much more glassy and crystalline than chalcedony
Hex.	C, none F, conchoidal, prominent	2.6	Geodes; botryoidal; banded; stalactitic	Geodes often have glassy quartz centers
Orth.	C, not important F, uneven Brittle	2.6	Massive; granular	Occurs in gneisses and schists with sillimanite, andalusite; resembles blue quartz
Hex.	C, rough basal F, uneven Brittle	2.6 2.8	Hexagonal prisms; broken crystals	Occurs in granite with quartz, feldspar, and mica
Orth.	C, basal, perfect and prominent Brittle	3.4 3.6	Crystals	Resembles aquamarine beryl except in crystal form; not common color
Hex.	C, rhombohedral, prominent Tough	3.9 4.1	Massive; grains; barrel-shaped crystals	Masses often show fine parallel striations due to twinning and cleavage
Orth.	C, basal, perfect and prominent F, splintery, uneven	2.8	Foliated massive	Soft and greasy feel; very flexible but not elastic
Mono.	C, clinopinacoidal, prominent in crystals Brittle	2.6	Long prisms with striated faces; earthy; powder	Earthy masses in clay, bones, fossils; crystals often in pyrrhotite

STREAK UNCOLORED, WHITE,

COLOR GREEN.	Name.	Composition.	Color.	Luster.	H.
	GARNIERITE	$H_2(Ni, Mg)SiO_4 + H_2O$	Apple-green	Dull	1 2
	CHLORITE PROCHLORITE CLINOCHLORE	$H_2(Mg, Fe)_2Al_2Si_2O_{10}$	Grass-green Brownish green Dark green	Pearly Vitreous	1.5 2.5
	MUSCOVITE (Chrome mica)	$H_2KAl_2(SiO_4)_2$ with Cr	Emerald-green Apple-green	Pearly Vitreous	2 2.5
	BIOTITE	$(HK)_2(Mg, Fe)_2(AlFe)_2(SiO_4)_2$	Brownish green Deep green	Pearly Vitreous	2.5 3
	CHALCANTHITE	$CuSO_4 + 5H_2O$	Bluish green	Vitreous Greasy	2.5
	CHRYSOCOLLA	$CuSiO_3 + 2H_2O$	Bluish green	Greasy Vitreous Dull	2 4
	SERPENTINE CHRYBOTILE ASBESTOS	$H_4Mg_2Si_2O_6$	Oil-green Light green Dark green Blackish green	Greasy Silky	2.5 4
	ACTINOLITE	$Ca(Mg, Fe)_3(SiO_3)_4$	Grass-green Deep green	Vitreous Silky	2.5 4
	BARITE	$BaSO_4$	Pale green	Vitreous Glassy	2.5 3.5
	WAVELLITE	$Al_2(OH)_2(PO_4)_2 + 5H_2O$	Pale green Bright green	Vitreous Pearly	3 4
	PYROMORPHITE	$(PbCl)Pb_4(PO_4)_3$	Yellowish green Dark green	Adamantine Pearly	3.5 4
	FLUORITE	CaF_2	Pale green Bright green Bluish green	Vitreous Glassy	4

System.	Cleavage or Fracture.	G.	Common Structure.	Observations.
Amorph.	C, none F, earthy	2.3 2.8	Friable masses; clay-like masses	Rounded and pod-shaped masses in clay
Mono.	C, basal, perfect and prominent Tough	2.8	Foliated; micaceous; scaly; flaky	Flexible but not elastic; dark-colored in thin plates; very common in schists
Mono.	C, basal, perfect and prominent Tough	2.7 3	Micaceous; scales; flakes; sheets	Light color to colorless in thin sheets; highly elastic
Mono.	C, basal, perfect and prominent Tough	2.7 3.1	Micaceous; scales; flakes	Dark-colored mica in thinnest sheets; elastic and flexible
Tric.	C, not prominent F, conchoidal Brittle	2.1 2.3	Crystals; massive; fibrous	Taste nauseous metallic; blue vitriol
Amorph.	C, none F, conchoidal Sectile to brittle	2 2.2	Incrustations; seams; stains	Never fibrous like malachite; often with malachite, chalcopyrite
Mono.	C, not important F, conchoidal or splintery	2.6	Massive; fibrous	Feels smooth and looks greasy; dark masses often intersected by veinlets of chrysotile asbestos
Mono.	C, fibrous Brittle	3	Fibrous reticulated masses	Occurs as actinolite schists; individual fibers are harder
Orth.	C, basal and prismatic, very prominent Brittle	4.3 4.6	Platy; massive; crystals	Usually nearly colorless with greenish cast; heavy vitreous mineral
Orth.	C, not prominent Brittle	2.3	Fine radiating fibrous globules; rosette-like	Usually on rock surface as small fibrous rosettes
Hex.	C, not prominent Brittle	6.5 7.1	Hexagonal prisms with striated faces; granular; fibrous	Often with galena, anglesite, mimetite
Isom.	C, octahedral, very prominent Brittle	3 3.2	Cubes; octahedral cleavage pieces; massive; granular	Often with calcite, galena, pyrite, dolomite

STREAK UNCOLORED, WHITE,

		Name.	Composition.	Color.	Luster.	H.	
COLOR GREEN.		CALAMINE	$H_2Zn_2SiO_5$	Bluish green Pale green	Vitreous	4.5 5	
		SMITHSONITE	$ZnCO_3$	Grayish green Bluish green	Vitreous	5	
		APATITE	$(CaF)Ca_4(PO_4)_3$	Pale green Grass-green Dark green Brownish green	Greasy Vitreous	5	
		OPAL	$SiO_2 + nH_2O$	Grayish green	Waxy Vitreous	5.5 6.5	
		WILLEMITE	Zn_2SiO_4	Yellowish green Bright green	Vitreous	5.5	
		CYANITE	Al_2SiO_5	Pale bluish green	Vitreous	5 7	
		PYROXENE	ENSTATITE	$MgSiO_3$	Grayish green Brownish green	Vitreous Pearly Silky	5.5
			DIOPSIDE	$CaMg(SiO_3)_2$	Pale green Bright green	Vitreous Glassy	5 6.5
			AUGITE	Silicate of Ca, Mg, Fe, and Al, chiefly	Blackish green	Vitreous	5 6
		AMPHIBOLE	ACTINOLITE	$Ca(Mg, Fe)_3(SiO_3)_4$	Grass-green Dark green	Vitreous Silky	5 6
			HORNBLLENDE	Silicate of Ca, Mg, Fe, and Al, chiefly	Blackish green	Vitreous Pearly	5 6
				TURQUOIS (Variscite)	$AlPO_4 \cdot Al(OH)_3 + H_2O$	Bluish green Apple-green	Waxy Dull

OR LIGHT GRAY.

System.	Cleavage or Fracture.	G.	Common Structure.	Observations.
Orth.	C, prismatic, sometimes prominent	3.5	Fibrous mammillary masses	Often with smithsonite
Hex.	C, rhombodehral, not usually prominent Brittle	4.4	Drusy masses; botryoidal or mammillary	Usually very compact, like chalcedony
Hex.	C, basal, not prominent Brittle	3.2	Hexagonal prisms; granular; massive	Commonly intermixed with brown colors; often with calcite; crystals often have fused appearance
Amorph.	C, none F, conchoidal and prominent	1.9 2.3	Colloidal masses	Waxy luster characteristic
Hex.	C, prismatic, not prominent Brittle	3.9 4.1	Massive; granular	Usually with franklinite, zincite, and calcite
Tric.	C, macropinacoidal, prominent Tough	3.6	Bladed; columnar	Divergent columnar; long blades usually bent and cross-fractured
Orth.	C, prismatic, prominent Brittle	3.1 3.3	Prismatic masses; divergent columns	Often much softer, owing to alteration to serpentine
Mono.	C, prismatic, not prominent Brittle	3.3	Crystals; square prisms with oblique base	Usually prisms have a prominent basal parting
Mono.	C, prismatic, not prominent Cleavage angle = 87°	3.3	Crystals; massive	Cleavage not so prominent as in hornblende; more common as crystals
Mono.	C, prismatic, prominent Cleavage angle = 124°	3 3.2	Divergent columnar or fibrous; reticulated masses	Often with talc or chlorite; fine to coarse fibrous and reticulated; often in schists
Mono.	C, prismatic and very prominent Cleavage angle = 124°	3 3.2	Massive; prismatic; columnar	Cleavage faces usually have fibrous appearance; common in granitic rocks and schists
None.	C, none Brittle	2.6	Globular masses; veins; seams	Usually intermixed with rock in irregular masses or veins

STREAK UNCOLORED, WHITE,

COLOR GREEN.	Name.	Composition.	Color.	Luster.	H.
	NEPHELITE (Elaeolite)	NaAlSiO_4	Grayish green Brownish green	Greasy Vitreous	5.5 6
	MICROCLINE (Feldspar)	KAlSi_3O_8	Bright green	Vitreous Pearly	6 6.5
	PREHNITE	$\text{H}_2\text{Ca}_2\text{Al}_2(\text{SiO}_4)_3$	Pale green Bright green	Vitreous	6 6.5
	CHLORITOID	$\text{H}_2(\text{Fe}, \text{Mg})\text{Al}_2\text{SiO}_7$	Dark green Greenish black	Pearly Vitreous	6.5
	EPIDOTE	$\text{HCa}_2(\text{Al}, \text{Fe})_3\text{Si}_3\text{O}_{13}$	Pistachio-green Yellowish green Oil-green Brownish green	Vitreous	6 7
	VESUVIANITE	$\text{MgCa}_3(\text{Al}, \text{OH})\text{Al}_2(\text{SiO}_4)_5$	Brownish green Bright green	Vitreous Greasy	6.5
	OLIVINE (Chrysolite)	$(\text{Mg}, \text{Fe})_2\text{SiO}_4$	Bottle-green Oil-green Grass-green	Vitreous Glassy	6.5 7
	JADEITE	$\text{NaAl}(\text{SiO}_3)_2$	Grayish green Deep green	Vitreous Silky	6.5 7
	QUARTZ	SiO_2	Light to dark green	Vitreous Glassy	7
	CHALCEDONY var. Jasper Chrysoprase Plasma	SiO_2	Apple-green Leek-green Light to dark green	Vitreous Waxy	7
	GARNET (Uvarovite)	$\text{Ca}_3\text{Cr}_2(\text{SiO}_4)_3$	Emerald-green	Vitreous	7 7.5
	TOURMALINE	$4\text{H}_2\text{O} \cdot 2(\text{Na}, \text{Li})_2\text{O} \cdot 3\text{B}_2\text{O}_3 \cdot 8\text{Al}_2\text{O}_3 \cdot 12\text{SiO}_2$	Dark green	Vitreous Glassy	7 7.5

System.	Cleavage or Fracture.	G.	Common Structure.	Observations.
Hex.	C, not prominent Brittle	2.6	Massive; short hexagonal prisms (rare)	Usually with augite, sodalite, leucite
Tric.	C, basal and brachypinacoidal Brittle	2.5	Crystals; cleavage pieces	Pearly luster on basal cleavage, also vein-like markings; green Amazon stone
Orth.	C, not prominent F, uneven Brittle	2.9	Reniform masses with drusy surfaces; small stalactitic	Occurs usually in cavities and fissures in basalts and diabases
Mono.	C, basal, prominent Brittle	3.5	Foliated massive; micaceous; platy	Deep green in thin plates; resembles hornblende
Mono.	C, basal, perfect and usually prominent Brittle	3.2 3.5	Crystals; prismatic; long columnar; granular; massive	Often in quartz; also with pyroxene, hornblende, magnetite, garnet
Tetrag.	C, not prominent F, uneven Brittle	3.4	Square prisms; faces often vertically striated; massive; granular	Often with calcite; crystals usually not perfect
Orth.	C, not important Brittle	3.3	Rounded masses of green grains; massive granular	Occurs in basalt as bomb-shaped masses of light and dark green, very glassy grains
Mono.	C, not prominent Tough	3.3	Very compact fibrous	Tough masses of interlocking fibers
Hex.	C, none F, conchoidal and prominent	2.6	Hexagonal prisms and pyramids	Quartz stained with chlorite or actinolite
Hex.	C, none F, conchoidal and prominent Brittle	2.6	Massive, compact; cryptocrystalline	Not so glassy as quartz
Isom.	C, none Brittle	3.5	Small crystals; granular	Sometimes on chromite as green glassy crystals
Hex.	C, none F, uneven Very brittle	3 3.2	Trigonal or hexagonal prisms	Often with pink tourmaline in lepidolite or with quartz, biotite, feldspar

STREAK UNCOLORED, WHITE,

	Name.	Composition.	Color.	Luster.	H.
COLOR GREEN.	BERYL var. Aquamarine Emerald Common	$\text{Be}_3\text{Al}_2(\text{SiO}_3)_6$	Pale green Bluish green Sea-green Emerald-green	Vitreous Very glassy	7.5 8
	TOPAZ	$\text{Al}_2(\text{F,OH})_2\text{SiO}_4$	Bluish green	Vitreous	8
	CHRYSOBERYL	BeAl_2O_4	Brownish green	Greasy Vitreous	8 8.5
	CORUNDUM	Al_2O_3	Bluish green Grayish green	Vitreous	9
COLOR BLACK.	BIOTITE (Mica)	$(\text{HK})_2(\text{Mg,Fe})_2(\text{Al,Fe})_2(\text{SiO}_3)_6$	Brownish black Greenish black	Vitreous Pearly	2.5 3
	CALCITE DOLOMITE (Limestone)	CaCO_3 $(\text{Ca,Mg})\text{CO}_3$	Grayish black	Vitreous	3 3.5 4
	FLUORITE	CaF_2	Dark purple- black	Vitreous	4
	HORNBLLENDE	Silicate of Ca, Mg, Fe, and Al, chiefly	Greenish black Brownish black	Vitreous Silky Pearly	5 6
	AUGITE	Silicate of Ca, Mg, Fe, and Al, chiefly	Greenish black Brownish black	Vitreous	5 6
	ALLANITE	$(\text{Ca,Fe})_2(\text{Al,Ce,Fe})_2(\text{AlOH})(\text{SiO}_3)_6$	Brownish black Pitch-black	Pitchlike Subme- tallie	5.5 6
	BROOKITE	TiO_3	Brownish black	Subme- tallie Adaman- tine	5.5 6

System.	Cleavage or Fracture.	G.	Common Structure.	Observations.
Hex.	C, rough basal, not prominent Brittle	2.6 2.8	Hexagonal prisms with basal planes; broken crystals	Often in granite with mica and feldspar
Orth.	C, basal, perfect and prominent Brittle	3.4 3.6	Crystals	Often nearly square prisms with base; resembles aquamarine beryl, except in form
Orth.	C, not prominent Brittle	3.5 3.8	Twinned crystals; tabular	Plates with twinning striations radiating from center; occurs with feldspar, garnet
Hex.	C, rhombohedral prominent Tough	3.9 4.1	Massive	Often with cleavage striations on faces
Mono.	C, basal, perfect and prominent Tough	2.7 3.1	Micaceous; plates; scales	Dark-colored mica in thinnest plates; common in granitic rocks
Hex.	C, none F, fine or coarse granular Brittle	2.7 2.9	Compact granular	Black limestone or marble
Isom.	C, octahedral, prominent Brittle	3 3.2	Massive; banded	Black color not common
Mono.	C, prismatic, prominent Cleavage angle 124°	2.9 3.4	Massive; fibrous; long prismatic	Cleavage faces very bright with often fibrous appearance; common with feldspar, quartz
Mono.	C, prismatic, not very prominent Cleavage angle 87°	3.3	Almost square prisms with oblique bases; massive	Usually with dark basaltic rocks; cleavage not so good as in hornblende
Mono.	C, not prominent F, uneven Brittle	3.5 4.2	Tabular crystals; thin seams in rock	Occurs in granitic rocks as black, pitch-like veins or as crystals
Orth.	C, not prominent Brittle	3.8 4.8	Crystals only; square or hexagonal-shaped pyramids	Crystal faces often striated; not twinned like rutile

STREAK UNCOLORED, WHITE,

	Name.	Composition.	Color.	Luster.	H.
COLOR BLACK.	RUTILE	TiO ₂	Brownish black	Metallic Adamantine	6 6.5
	CASSITERITE	SnO ₂	Black	Submetallic Adamantine	6 7
	QUARTZ	SiO ₂	Grayish black Brownish black	Vitreous	7
	GARNET var. Melanite	Silicate of Ca, Fe, Al, and Ti	Velvet-black Brownish black	Vitreous Velvety	7 7.5
	TOURMALINE	Borosilicate of Al, Fe, and Mg	Coal-black	Vitreous Very glassy	7 7.5
	SPINEL	(Mg,Fe)Al ₂ O ₄	Grayish black	Vitreous Dull	8
COLOR WHITE, GRAY, OR COLORLESS.	ULEXITE	NaCaB ₆ O ₆ +8H ₂ O	Snow-white	Pearly Silky	1
	TALC Soapstone	H ₂ Mg ₂ (SiO ₃) ₄	White Greenish white Gray	Pearly Greasy Dull	1 1.5
	PYROPHYLLITE	HA1(SiO ₃) ₂	White Grayish Brownish gray	Pearly Greasy Dull	1 2
	CERARGYRITE (Hornsilver)	AgCl	Gray Brownish gray	Resinous Waxy	1 1.5
	SAL AMMONIAC	NH ₄ Cl	White Gray Colorless	Vitreous	1.5 2
	CALCITE var. Chalk	CaCO ₃	White	Earthy Dull	1.5 2.5

System.	Cleavage or Fracture.	G.	Common Structure.	Observations.
Tetrag.	C, not prominent Brittle	4 2	Crystals, usually twinned and faces deeply striated	Crystals generally imperfect; knee-shaped twins common
Tetrag.	C, not prominent F, uneven, coarse Brittle	6 8 7.1	Massive; granular; rolled pebbles; twinned crystals	Often with quartz, mica, wolframite, fluorite; heavy black masses
Hex.	C, none F, conchoidal Brittle	2 6	Crystals; hexagonal prisms and pyramids	Very dark smoky quartz
Isom.	C, none F, uneven Brittle	3 8	Crystals; rhombic dodecahedrons	Uncommon color
Hex.	C, none F, uneven Very brittle	3 3.2	Crystals; long trigonal-shaped prisms; sometimes divergent columnar	Crystal faces usually striated vertically, and much fractured horizontally; often as coal-black crystals in quartz and feldspar
Isom.	C, imperfect F, conchoidal Brittle	3.5 4 1	Crystals; octahedrons	In granular limestone often with chondrodite
	C, not important F, fibrous	1 6	Soft fibrous masses	Usually in ball like masses of fibers
Mono.	C, basal, perfect and prominent in the foliated masses	2 8	Foliated; compact massive; fibrous	Soft and greasy feel; fibers usually not radiate like pyrophyllite
Mono	C, basal and prominent Flexible	2.9	Fibrous, radiate; foliated; massive	Often in small hemispheres of radiating fibers; soft and greasy like talc
Isom.	C, none Sectile	5.5	Wax-like crusts; horn-like masses	Cuts like wax; often with ores of silver
Isom.	C, not important Brittle	1.5	Crusts; globular masses	Occurs on lava rock; disagreeable saline taste
	C, none Brittle	2.7	Soft white earthy masses	Resembles white kaolinite, but has no clay odor

Name.	Composition.	Color.	Luster.	H.
GYPSUM var. Selenite Alabaster Satin-spar Common	$\text{CaSO}_4 + 2\text{H}_2\text{O}$	Colorless White Gray	Pearly Vitreous Silky Dull	1.5 2
KAOLINITE (Clay)	$\text{H}_2\text{Al}_2\text{Si}_2\text{O}_7$	White Gray Colorless	Dull Earthy Greasy	2 2.5
BAUXITE	$\text{Al}_2\text{O}_3 + 3\text{H}_2\text{O}$	White Gray	Earthy Dull	2 2.5
SEPIOLITE (Meerschaum)	$\text{H}_4\text{Mg}_2\text{Si}_5\text{O}_{16}$	White	Earthy Dull	2 2.5
BORAX	$\text{Na}_2\text{B}_4\text{O}_7 + \text{H}_2\text{O}$	Snow-white Colorless	Earthy Dull Vitreous	2 2.5
KALINITE (Alum)	$\text{AlK}(\text{SO}_4)_2 + 12\text{H}_2\text{O}$	White Colorless	Vitreous Icy	2.5
EPSOMITE	$\text{MgSO}_4 + 7\text{H}_2\text{O}$	White	Vitreous	2 2.5
HALITE	NaCl	Colorless White Bluish white	Vitreous	2.5
BRUCITE	$\text{Mg}(\text{OH})_2$	White Greenish white	Pearly	2.5
TREMOLITE var. Asbestos Mountain leather Mountain cork	$\text{CaMg}_3(\text{SiO}_3)_4$	White Gray	Silky Pearly	2 2.5
SERPENTINE var. Chrysotile or Asbestos	$\text{H}_3\text{Mg}_3\text{Si}_2\text{O}_7$	Greenish white	Silky	2.5 4

COLOR WHITE, GRAY, OR COLORLESS.

System.	Cleavage or Fracture.	G.	Common Structure.	Observations.
Mono.	C, clinopinacoidal, prominent in selenite Sectile	2.3	Colorless crystals and cleavage plates; massive white; fibrous; columnar; granular	Soft and brittle; not so pearly luster as brucite, and softer; compact translucent masses common
Mono.	C, basal in crystals, none in massive F, earthy Brittle or sectile	2.6	Compact massive; rarely in colorless flakes	Rough feeling; soapy var has greasy feel; strong odor of clay when breathed on
	C, none F, earthy Brittle	2.5	Compact massive; pisolitic	Distinguished from clay only by pea-shaped structure
Mono.	C, none F, earthy	2	Massive; mammillary; reniform; very compact	Very smooth feel; has not the clay odor of kaolinite
Mono.	C, orthopinacoidal, not prominent Brittle	1.7	Crystals; powder	Taste alkaline; white crystals often have fresh, unaltered glassy centers
Isom.	C, none Brittle	1.7	Crystals; octahedrons; mealy crusts	Alum taste
Orth.	C, brachypinacoidal, prominent Brittle	1.7	Long acicular crystals; capillary tufts; efflorescences	Taste bitter and salt; often in sulphide mines as efflorescences on walls
Isom.	C, cubic, perfect and prominent Brittle	2.1 2.6	Cubes; massive; granular	Salt taste; sometimes with anhydrite
Hex.	C, basal, perfect and prominent Flexible	2.5	Foliated; massive	Resembles gypsum but has more pearly luster; often with serpentine
Mono.	C, fibrous F, fibrous Brittle	<1 3	Fibrous; asbestiform; sheets; cork-like masses	Occurs with tremolite, feldspar quartz; not green like chrysotile when compact
Mono.	C, fibrous Brittle	2.6	Fibrous; asbestiform	Narrow fibrous veins in serpentine; fibers are green in compact mass

STREAK UNCOLORED, WHITE

COLOR WHITE, GRAY, OR COLORLESS.	Name.	Composition.	Color.	Luster.	H.
	ANDALUSITE var. Chiastolite	Al_2SiO_5	Dark gray Blackish gray'	Vitreous	2 4
	MUSCOVITE (Mica)	$H_2KAl_3(SiO_4)_3$	Colorless Gray	Pearly Vitreous	2 2.5
	LEPIDOLITE (Mica)	$(LiK)_2Al_2(F,OH)_2Si_2O_6$	Pale pinkish white Lavender Gray	Pearly	2.5 4
	CRYOLITE	Na_3AlF_6	Pure white	Icy Vitreous	2.5
	CALCITE var. Iceland spar Stalactites Marble Common	$CaCO_3$	White Gray Colorless	Vitreous Glassy	3
	ANGLESITE	$PbSO_4$	Gray White Colorless	Adaman- tine Greasy Dull	3
	CERUSSITE	$PbCO_3$	Cream-white Gray	Adaman- tine	3 3.5
	BARITE	$BaSO_4$	White Colorless Gray Yellowish white	Vitreous Pearly	2.5 3.5
	ANHYDRITE	$CaSO_4$	White Bluish white Reddish white Gray	Vitreous Pearly	3 3.5
CELESTITE	$SrSO_4$	Colorless with bluish tinge White	Vitreous Glassy	3 3.5	

OR LIGHT GRAY.

System.	Cleavage or Fracture.	G.	Common Structure.	Observations.
Orth.	C, not prominent Brittle	3.2	Rounded prisms; square prisms	Occurs in schists often as knotty projections; end sections show black squares or crosses
Mono.	C, basal, perfect and very prominent Tough	2.7 3	Micaceous; large sheets; foliated; flakes; scales	Highly flexible and elastic; colorless in thin sheets; cleavable in the thinnest sheets
Mono.	C, basal, perfect and prominent Tough	2.9	Fine or coarse scaly masses; platy; micaceous; foliated	Compact scaly masses containing pink tourmaline
Mono.	C, basal and pinacoidal; basal is prominent	3	Massive	Snow-ice appearance; often with siderite; cleavage in three directions almost at right angles
Hex.	C, rhombohedral, very perfect and prominent Brittle	2.7	Crystals; rhombohedrons; scalenohedrons; granular; stalactitic; banded, etc.	Commonly associated with the metallic minerals; colorless variety is Iceland spar; calcite is apt to be stained any color
Orth.	C, not prominent F, conchoidal Brittle	6.1 6.3	Massive, often banded; crystals	Occurs with galena as an alteration product; crystals are colorless; gray masses often have core of galena
Orth.	C, not prominent Very brittle	6.5	Prismatic crystals; massive	Occurs similar to angle-site; gray masses somewhat porous or reticulated
Orth.	C, basal and prismatic, prominent	4.3 4.6	Crystals; crested masses; granular; lamellar; concretions massive	Often with galena; heavy white mineral, called heavy spar
Orth.	C, pinacoidal, prominent Brittle	3	Massive; granular; scaly	Cleavage in three directions at right angles, making cube forms, occurs with gypsum, limestone
Orth.	C, basal and prismatic; basal very prominent	3.9	Cleavage masses; crystals	Often as colorless crystals with native sulphur

STREAK UNCOLORED, WHITE,

Name.	Composition.	Color.	Luster.	H.
WITHERITE	BaCO_3	White	Vitreous	3 3.5
STRONTIANITE	SrCO_3	White Yellowish white	Vitreous Glassy	3 3.5
ARAGONITE	CaCO_3	White Gray Colorless	Vitreous Glassy	3.5 4
DOLOMITE	$(\text{CaMg})\text{CO}_3$	White Gray	Vitreous	3.5 4
SIDERITE	FeCO_3	Brownish gray	Vitreous Pearly	3.5 4
FLUORITE	CaF_2	Greenish white White Colorless	Vitreous Glassy	4
COLEMANITE	$\text{Ca}_2\text{B}_6\text{O}_{11} + 5\text{H}_2\text{O}$	Colorless White Yellowish white	Vitreous Very glassy	4 4.5
SCHEELITE	CaWO_4	Gray Yellowish	Adamantine Greasy	4.5 5
WOLLASTONITE	CaSiO_3	White Gray	Vitreous	4.5 5
CHABAZITE	$\text{Ca}_2\text{Al}_2(\text{SiO}_4)_2(\text{Si}_2\text{O}_7)_2 + 18\text{H}_2\text{O}$	White Colorless Gray	Vitreous	4 5
APOPHYLLITE	$\text{H}_7\text{KCa}_4(\text{SiO}_3)_8 + 4\frac{1}{2}\text{H}_2\text{O}$	White Colorless Yellowish	Vitreous Glassy Pearly on base	4.5 5


COLOR WHITE, GRAY, OR COLORLESS.

System.	Cleavage or Fracture.	G.	Common Structure.	Observations.
Orth.	C, not prominent Brittle	4.3	Columnar; hexagonal-shaped crystals with striated faces	Sometimes with galena; heavy snow-white masses common
Orth.	C, prismatic, sometimes prominent Brittle	3.7	Columnar masses	Divergent columnar masses resembling aragonite or calcite, but much heavier
Orth.	C, prismatic but not usually prominent Brittle	2.9	Stalactitic; banded; columnar; hexagonal-shaped crystals	Distinguished from calcite by lack of cleavage and by hardness
Hex.	C, rhombohedral, sometimes prominent Brittle	2.9	Rhombohedrons with curved faces; massive; granular	Massive variety indistinguishable from calcite except somewhat harder; crystals have curved faces
Hex.	C, rhombohedral, very prominent Brittle	3.8	Rhombohedrons with curved faces; saddle-shaped masses; compact; massive	Darker and heavier than dolomite; often as rhombohedrons in cryolite
Isom.	C, octahedral, very prominent Brittle	3 3.2	Cubes; octahedrons; massive; granular	Often with magnetite, pyrite, calcite; sometimes very compact granular
Mono.	C, clinopinacoidal, very prominent Brittle	2.4	Crystals; massive	Cleaves into thin brittle plates
Tetrag.	C, not prominent Brittle	5.9 6.1	Crystals; pyramids; massive	Often with cassiterite, wolframite, purple fluorite; very heavy
Mono.	C, orthopinacoidal, not prominent Brittle	2.9	Fibrous; columnar	Parallel, or reticulated, fibrous masses; often in marble; resembles tremolite
Hex.	C, not prominent Brittle	2.1	Crystals, almost cubic in shape	Usually in cavities of lava rock with stilbite, heulandite, natrolite
Tetrag.	C, basal, perfect and prominent Brittle	2.4	Crystals; short prisms with base; also pointed pyramids	Basal cleavage has very pearly luster, prismatic faces glassy and vertically striated

STREAK UNCOLORED, WHITE,

COLOR WHITE, GRAY OR COLORLESS.	Name.	Composition.	Color.	Luster.	H.
	CALAMINE	$H_2Zn_2SiO_4$	Colorless White Gray	Vitreous	4.5 5
	MAGNESITE	$MgCO_3$	Snow-white Gray	Vitreous Dull	3.5 4.5
	SMITHSONITE	$ZnCO_3$	Bluish gray Yellowish gray	Vitreous	5
	APATITE	$(CaF)Ca_3(PO_4)_3$	Colorless Gray	Vitreous Greasy	5
	PECTOLITE	$HNaCa_2(SiO_3)_2$	White	Silky Vitreous	5
	NATROLITE	$H_4NaAl_2(SiO_3)_4$	White Colorless	Vitreous Silky	5 5.5
	DATOLITE	$H_4Ca(BO)SiO_4$	Colorless White	Vitreous Glassy	5 5.5
	ANALCITE	$Na_2Al_2(SiO_3)_4 \cdot 2H_2O$	Colorless White	Vitreous Glassy	5 5.5
	OPAL	$SiO_2 + nH_2O$	Gray White	Waxy Vitreous	5.5 6.5
SCAPOLITE WERNERITE	Silicate of Ca, Al, Na, and Cl	Gray Greenish gray White	Vitreous Silky	5.5 6	
LEUCITE	$KAl(SiO_3)_2$	Gray White	Vitreous	5.5 6	

System.	Cleavage or Fracture.	G.	Common Structure.	Observations.
Orth.	C, prismatic and prominent Brittle	3.5	Drusy coatings and crusts; small crystals	Often on yellowish brown earthy masses, in small drusy crystallizations
Hex.	C, rhombohedral in crystals F, conchoidal and prominent	3.1	Crystals rare; massive, impalpable	Very compact tough white masses with soiled surfaces; are apparently very hard
Hex.	C, rhombohedral, but not usually prominent Brittle	4.3 4.5	Botryoidal crusts; drusy crusts	Often with sphalerite or calamine
Hex.	C, basal, not prominent Brittle	3.2	Crystals; massive	Common white phosphate rock; crystals are colorless, green or brown usually
Mono.	C, not prominent Brittle to tough	2.7	Fibrous; divergent, radiate, reticulated	Long white fibers divergent to sharp points; also compact fibrous
Orth.	C, prismatic, prominent in coarse varieties Brittle	2.2	Acicular; coarse; columnar; fibrous	Often with stilbite, apophyllite, analcite, chabazite, in cavities of lava rock
Mono.	C, none F, uneven Brittle	3	Crystals; massive; granular to compact	Small glassy crystals with slight greenish tint on lava rock; massive white
Isom.	C, not prominent Brittle	2.2	Crystals; icosatetrahedrons or cubes	Often in cavities of lava with apophyllite, natrolite, chabazite, prehnite, datolite
Amorph.	C, none F, conchoidal and very prominent	1.9 2.3	Massive; colloidal; blebby; globular	Wood opal; common opal
Tetrag.	C, not prominent Brittle	2.5 2.8	Square prisms with low pyramidal ends; massive	Crystals usually have rough, uneven faces; often in crystalline limestone
Tetrag.	C, imperfect F, conchoidal Brittle	2.5	Crystals; trapezohedrons	Always in crystals; occurs in volcanic rocks with nephelite, sodalite



STREAK UNCOLORED, WHITE,

	Name.	Composition.	Color.	Luster.	H.	
COLOR WHITE, GRAY, OR COLORLESS.	ENSTATITE	$MgSiO_3$	Greenish gray	Pearly Vitreous	5.5	
	PYROXENE DIOPSIDE	$CaMg(SiO_3)_2$	Colorless Yellowish white Greenish white	Glassy Vitreous	6 6.5	
	TREMOLITE	$CaMg_3(SiO_3)_4$	White Gray	Silky Pearly Vitreous	5 6	
	NEPHELINE ELÆOLITE	$NaAlSi_3O_8$	Greenish gray Brownish gray	Greasy Vitreous	5.5 6	
	AMBLYGONITE	$Li(Al,F)PO_4$	White	Vitreous	6	
	FELDSPARS	ORTHOCLASE	$KAlSi_3O_8$	White Gray Colorless	Vitreous Pearly	6 6.5
		MICROCLINE	$KAlSi_3O_8$	White Gray Yellowish	Vitreous Pearly	6 6.5
		ALBITE	$NaAlSi_3O_8$	White Colorless Gray	Glassy Vitreous	6 6.5
		OLIGOCLASE	$NaAlSi_3O_8 + CaAl_2Si_2O_8$	Colorless White	Vitreous Glassy	6 6.5
		LABRADORITE	$CaAl_2Si_2O_8 + NaAlSi_3O_8$	Dark gray Grayish white	Vitreous Pearly	5 6
		ANORTHITE	$CaAl_2Si_2O_8$	White Gray	Vitreous	6 6.5

OR LIGHT GRAY.

67

System.	Cleavage or Fracture.	G.	Common Structure.	Observations.
Orth.	C, prismatic and pinacoidal, prominent Brittle	3.1 3.3	Columnar, divergent masses; lamellar	Often soft owing to alteration to serpentine; cleavage faces quite pearly or silky in luster
Mono.	C, prismatic, not prominent Brittle	3.3	Crystals, almost square or rounded	Often with blue calcite, brown tourmaline
Mono.	C, prismatic, prominent Cleavage angle 124°	2.9 3.1	Columnar; fibrous; prismatic crystals	Often as crystals in dolomitic limestone or marble; also as compact fibrous masses
Hex.	C, not prominent Brittle	2.6	Massive; sometimes hexagonal prisms	Often with sodalite, albite, leucite; greasy luster characteristic
Tric.	C, basal, perfect and prominent Brittle	3	Compact massive	Often with lepidolite, tourmaline
Mono.	C, basal and clinopinacoidal, prominent Brittle	2.4 2.6	Crystals; massive; cleavage pieces	Two cleavages at right angles; common in granitic rocks with mica, hornblende, and quartz
Tric.	C, basal and brachypinacoidal, prominent	2.5	Crystals; massive	Usually has fine crossveined structure on the basal plane
Tric.	C, basal and brachypinacoidal, not so prominent	2.6	Small crystals; twinned crystals; platy masses	Fine parallel striations or reentrant angles on the base due to twinning
Tric.	C, basal and brachypinacoidal, not so prominent	2.6	Crystals; massive	Fine parallel striations on the basal cleavage due to twinning
Tric.	C, basal and brachypinacoidal, prominent	2.7	Massive; cleavage pieces	Fine striations on basal cleavage due to twinning; usually shows a beautiful play of colors; blue, green, gold, etc.
Tric.	C, basal, prominent F, uneven Brittle	2.7	Crystals; prismatic	Occurs in volcanic lavas; not so common as the other feldspars

STREAK UNCOLORED, WHITE,

COLOR WHITE, GRAY OR COLORLESS.	Name.	Composition.	Color.	Luster.	H.
	ZOISITE	$\text{Ca}_2\text{Al}_2(\text{AlOH})(\text{SiO}_3)_2$	Grayish white Greenish gray	Vitreous Pearly	6 6.5
	SPODUMENE	$\text{LiAl}(\text{SiO}_3)_2$	Gray White	Vitreous Pearly	6.5 7
	DIASPORE	$\text{AlO}(\text{OH})$	Lavender-gray Grayish white Cream white	Pearly Vitreous Adamantine	6.5 7
	QUARTZ var. Rock crystal Milky Smoky Common	SiO_2	Colorless White Smoky gray	Vitreous Greasy	7
	CHALCEDONY var. Agate Chert Flint Hornstone Siliceous sinter	SiO_2	Gray White	Waxy Vitreous	7
	ANDALUSITE (CHIASTOLITE)	Al_2SiO_5	Gray Reddish gray	Vitreous	7.5
	LAWSONITE	$\text{H}_4\text{CaAl}_2\text{Si}_2\text{O}_{10}$	Bluish white Gray	Vitreous	7.5 8
	ZIRCON	ZrSiO_4	Brownish gray Lavender-gray Colorless	Vitreous Pearly	7.5 8
	TOPAZ	$\text{Al}_2(\text{F},\text{OH})_2\text{SiO}_4$	White Colorless	Vitreous Glassy	8
CORUNDUM	Al_2O_3	Gray Bluish gray Greenish gray	Vitreous	9	
DIAMOND	C	Colorless Gray Yellowish	Adamantine	10	

*stains blue green
d.b. red
root*

OR LIGHT GRAF

System.	Cleavage or Fracture.	G.	Common Structure.	Observations.
Orth.	C, brachypinacoidal prominent Brittle ¹	3.3	Columnar; fibrous; prismatic	Columns deeply striated vertically, and broken transversely
Mono.	C, prismatic often prominent Brittle	3.2	Large crystals and broad cleavage plates lamellar	Often parts in broad lamellar masses
Orth.	C, brachypinacoidal; prominent Brittle	3.3 3.5	Bladed; foliated	Often as veins in chlorite with margarite, corundum; very pearly to adamantine luster
Hex.	C, none F, conchoidal and prominent Brittle	2.6	Hexagonal prisms and pyramids; massive; granular; sand; pebbles	Commonest mineral; occurs in most crystalline rocks as a constituent; conchoidal fracture is characteristic
Hex.	C, none F conchoidal and prominent Brittle to tough	2.6	Geodal; botryoidal; mammillary; banded; concretionary; massive	Not glassy like quartz; cryptocrystalline in structure; banded varieties classed as agates; geodes often have quartz centers
Orth.	C, not prominent F, uneven Brittle	3.2	Almost square prisms with broken ends	End sections of chiasolite show black crosses or squares due to inclusions
Orth.	C, brachypinacoidal; prominent Brittle	3.1	Crystals; lenticular plates	Often with margarite, actinolite, chlorite; resembles corundum
Tetrag.	C, none Brittle	4.7	Crystals; prisms and pyramids	Occurs in granites and syenites; loose crystals in gold sands
Orth.	C, basal, perfect and prominent Brittle	3.4 3.6	Crystals; white massive	Massive white distinguished from white quartz by presence of cleavage faces
Hex.	C, rhombohedral, prominent Tough	3.9 4.1	Massive; barrel-shaped crystals	Often with chlorite, margarite, magnetite; massive; has usually fine parallel parting striations
Isom.	C, octahedral, not prominent Brittle	3.5	Small rounded octahedral-shaped crystals	Occurs in dark bluish-green igneous rock

INDEX.

	PAGE		PAGE
Actinolite.....	48, 50	Brookite.....	24, 54
Agate.....	36, 42, 68	Brucite.....	58
Alabandite.....	28		
Alabaster.....	58	Calamine.....	32, 44, 50, 64
Albite.....	66	Calaverite.....	16
Allanite.....	36, 54	Calcite.....	30, 38, 44, 54, 56, 60
Almandite.....	38, 42	Carnelian.....	42
Alum.....	58	Cassiterite.....	22, 26, 36, 56
Amblygonite.....	66	Celestite.....	44, 60
Amethyst.....	42	Cerargyrite.....	28, 56
Amphibole.....	34, 50	Cerussite.....	30, 60
Analcite.....	64	Chabazite.....	40, 62
Andalusite.....	42, 60, 68	Chalcanthite.....	44, 48
Andradite.....	38, 42	Chalcedony.....	36, 42, 46, 52, 68
Anglesite.....	60	Chalcocite.....	10
Anhydrite.....	60	Chalcopyrite.....	14
Annabergite.....	26	Chalk.....	56
Anorthite.....	66	Chert.....	68
Anthophyllite.....	34	Chiasolite.....	60, 68
Antimony.....	12, 16	Chloanthite.....	14
Apatite.....	32, 50, 64	Chlorite.....	26, 30, 48
Apophyllite.....	40, 62	Chloritoid.....	52
Aquamarine.....	54	Chondrodite.....	36, 42
Aragonite.....	32, 62	Chromite.....	20, 24
Argentite.....	10	Chrysoberyl.....	54
Arsenic.....	12, 16	Chrysocolla.....	26, 44, 48
Arsenopyrite.....	14	Chrysolite.....	52
Asbestos.....	48, 58	Chrysoprase.....	52
Augite.....	28, 34, 50, 54	Chrysotile.....	48, 58
Axinite.....	36	Cinnabar.....	18
Azurite.....	26	Clay.....	28, 58
		Clinochlore.....	26, 30, 48
Barite.....	30, 44, 48, 60	Cobaltite.....	14
Bauxite.....	18, 24, 28, 58	Colemanite.....	62
Beryl.....	38, 46, 54	Columbite.....	12, 22
Biotite.....	30, 48, 54	Copper.....	18
Bismuth.....	16	Cordierite.....	46
Borax.....	58	Corundum.....	44, 46, 54, 68
Bornite.....	14	Cryolite.....	30, 60
Bronzite.....	34	Cuprite.....	18

	PAGE
Cyanite.....	46, 50
Datolite.....	64
Diamond.....	68
Diaspore.....	68
Diopside.....	50, 66
Dolomite.....	82, 40, 54, 62
Elæolite.....	36, 52, 66
Emerald.....	54
Enargite.....	10, 14
Enstatite.....	34, 50, 66
Epidote.....	26, 36, 52
Epsomite.....	58
Erythrite.....	18
Essonite.....	42
Famatinite.....	14
Feldspar.....	52, 66
Fibrolite.....	36
Flint.....	36, 68
Fluorite.....	32, 40, 44, 48, 54, 62
Franklinite.....	12, 20
Galenite.....	10, 12
Garnet.....	38, 42, 52, 56
Garnierite.....	48
Glaucophane.....	28
Gold.....	22
Göthite or Goethite.....	24
Graphite.....	10
Grossularite.....	38, 42
Gypsum.....	28, 38, 58
Halite.....	30, 58
Hematite.....	18, 20
Heulandite.....	40
Hornblende.....	28, 34, 50, 54
Hornsilver.....	56
Hornstone.....	68
Hypersthene.....	34
Iceland spar.....	60
Ilmenite.....	12, 20
Iolite.....	46
Iron.....	12
Jadeite.....	52
Jamesonite.....	10
Jasper.....	36, 42, 52
Kalinite.....	58
Kaolinite.....	28, 58

	PAGE
Labradorite.....	66
Lapis Lazuli.....	28, 44
Lazulite.....	44
Lazurite.....	28, 44
Lawsonite.....	68
Lepidolite.....	38, 60
Leucite.....	64
Limestone.....	54
Limonite.....	22, 24
Linarite.....	26
Magnesite.....	32, 64
Magnetite.....	12
Malachite.....	26
Manganite.....	20
Marble.....	60
Marcasite.....	14, 16
Margarite.....	40
Meerschäum.....	58
Melanite.....	56
Menaccanite.....	12
Mercury.....	16
Mica.....	60
Microcline.....	52, 66
Millerite.....	14
Mimetite.....	30
Molybdenite.....	10, 16
Monazite.....	34
Mountain cork.....	58
Mountain leather.....	58
Muscovite.....	48, 60
Natrolite.....	64
Nephelite.....	36, 52, 66
Nicolite.....	16
Oligoclase.....	66
Olivine.....	52
Opal.....	34, 40, 46, 50, 64
Orpiment.....	22
Orthoclase.....	36, 42, 66
Pectolite.....	64
Penninite.....	30
Phlogopite.....	30
Plasma.....	52
Platiniridium.....	16
Platinum.....	16
Prehnite.....	52
Prochlorite.....	26, 30, 48
Proustite.....	18
Psilomelane.....	12, 20

	PAGE		PAGE
Pyrargyrite	20	Stibnite	10, 12
Pyrite	16	Stilbite	32
Pyrolusite	10	Strontianite	62
Pyromorphite	24, 26, 30, 48	Sulphur	22, 28
Pyrope	38, 42	Sylvanite	16
Pyrophyllite	56		
Pyroxene	34, 50, 66	Talc	46, 56
Pyrrhotite	14	Tennantite	12
Quartz	36, 42, 46, 52, 56, 68	Tetrahedrite	12, 20
Realgar	22	Thulite	42
Rhodochrosite	40	Titanite	34
Rhodonite	40	Topaz	38, 44, 46, 54, 68
Rutile	24, 36, 42, 56	Tourmaline	38, 42, 52, 56
Sal Ammoniac	56	Tremolite	34, 58, 66
Satin Spar	58	Troostite	34
Scapolite	40, 64	Turgite	18
Scheelite	32, 62	Turquoise	46, 50
Selenite	58	Ulexite	56
Sepiolite	58	Uvarovite	52
Serpentine	30, 48, 58	Vanadinite	38
Siderite	24, 32, 62	Variscite	50
Siliceous Sinter	68	Vesuvianite	38, 52
Sillimanite	36	Vivianite	26, 44, 46
Silver	16	Wad	18
Smaltite	14	Wavellite	48
Smithsonite	32, 50, 64	Wernerite	40, 64
Soapstone	56	Willemite	34, 50
Sodalite	46	Witherite	62
Spessartite	38, 42	Wolframite	20
Sphalerite	18, 20, 22, 24, 32, 40	Wollastonite	62
Spinel	42, 56	Wulfenite	30, 38
Spodumene	68		
Stalactite	60	Zincite	22
Staurolite	38, 42	Zircon	38, 68
Stephanite	10	Zoisite	36, 42, 68

3) Gypsum

9) Anhydrite

14) Calcite

23) Cryscolla

28) Apatite



THIS BOOK IS DUE ON THE LAST DATE
STAMPED BELOW

AN INITIAL FINE OF 25 CENTS
WILL BE ASSESSED FOR FAILURE TO RETURN
THIS BOOK ON THE DATE DUE. THE PENALTY
WILL INCREASE TO 50 CENTS ON THE FOURTH
DAY AND TO \$1.00 ON THE SEVENTH DAY
OVERDUE.

SEP 2 1932

NOV 13 1935

NOV 27 1935

DEC 11 1935

MAR 31 1947

FEB 7 1955

DEC 6 1954

REG. CH. MAR 12 1979

INTERLIBRARY LOAN

F/R/R
JAN 5 1979

UNIV. OF CALIF., BERK.

YC 15051

119595
984m
E II

